

Research Article

Solution and Stability of a Mixed Type Additive, Quadratic, and Cubic Functional Equation

M. Eshaghi Gordji,¹ S. Kaboli Gharetapeh,² J. M. Rassias,³ and S. Zolfaghari¹

¹ Department of Mathematics, Semnan University, P.O. Box 35195-363, Semnan, Iran

² Department of Mathematics, Payame Noor University of Mashhad, Mashhad, Iran

³ Section of Mathematics and Informatics, Pedagogical Department, National and Capodistrian University of Athens, 4 Agamemnonos St., Aghia Paraskevi, Athens 15342, Greece

Correspondence should be addressed to M. Eshaghi Gordji, madjid.eshaghi@gmail.com

Received 24 January 2009; Revised 13 April 2009; Accepted 26 June 2009

Recommended by Patricia J. Y. Wong

We obtain the general solution and the generalized Hyers-Ulam-Rassias stability of the mixed type additive, quadratic, and cubic functional equation $f(x + 2y) - f(x - 2y) = 2(f(x + y) - f(x - y)) + 2f(3y) - 6f(2y) + 6f(y)$.

Copyright © 2009 M. Eshaghi Gordji et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

The stability problem of functional equations originated from a question of Ulam [1] in 1940, concerning the stability of group homomorphisms. Let (G_1, \cdot) be a group, and let $(G_2, *)$ be a metric group with the metric $d(\cdot, \cdot)$. Given $\epsilon > 0$, does there exist a $\delta > 0$, such that if a mapping $h : G_1 \rightarrow G_2$ satisfies the inequality $d(h(x \cdot y), h(x) * h(y)) < \delta$ for all $x, y \in G_1$, then there exists a homomorphism $H : G_1 \rightarrow G_2$ with $d(h(x), H(x)) < \epsilon$ for all $x \in G_1$? In other words, under what condition does there exist a homomorphism near an approximate homomorphism?

In 1941, Hyers [2] gave a first affirmative answer to the question of Ulam for Banach spaces. Let $f : E \rightarrow E'$ be a mapping between Banach spaces such that

$$\|f(x + y) - f(x) - f(y)\| \leq \delta, \quad (1.1)$$

for all $x, y \in E$ and for some $\delta > 0$. Then there exists a unique additive mapping $T : E \rightarrow E'$ such that

$$\|f(x) - T(x)\| \leq \delta, \quad (1.2)$$

for all $x \in E$. Moreover if $f(tx)$ is continuous in t for each fixed $x \in E$, then T is linear (see also [3]). In 1950, Aoki [4] generalized Hyers' theorem for approximately additive mappings. In 1978, Th. M. Rassias [5] provided a generalization of Hyers' theorem which allows the Cauchy difference to be unbounded. This new concept is known as Hyers-Ulam-Rassias stability of functional equations (see [2–24]).

The functional equation

$$f(x+y) + f(x-y) = 2f(x) + 2f(y) \quad (1.3)$$

is related to symmetric biadditive function. In the real case it has $f(x) = x^2$ among its solutions. Thus, it has been called quadratic functional equation, and each of its solutions is said to be a quadratic function. Hyers-Ulam-Rassias stability for the quadratic functional equation (1.3) was proved by Skof for functions $f : A \rightarrow B$, where A is normed space and B Banach space (see [25–28]).

The following cubic functional equation was introduced by the third author of this paper, J. M. Rassias [29, 30] (in 2000-2001):

$$f(x+2y) + 3f(x) = 3f(x+y) + f(x-y) + 6f(y). \quad (1.4)$$

Jun and Kim [13] introduced the following cubic functional equation:

$$f(2x+y) + f(2x-y) = 2f(x+y) + 2f(x-y) + 12f(x), \quad (1.5)$$

and they established the general solution and the generalized Hyers-Ulam-Rassias stability for the functional equation (1.5).

The function $f(x) = x^3$ satisfies the functional equation (1.5), which explains why it is called cubic functional equation.

Jun and Kim proved that a function f between real vector spaces X and Y is a solution of (1.5) if and only if there exists a unique function $C : X \times X \times X \rightarrow Y$ such that $f(x) = C(x, x, x)$ for all $x \in X$, and C is symmetric for each fixed one variable and is additive for fixed two variables (see also [31–33]).

We deal with the following functional equation deriving from additive, cubic and quadratic functions:

$$f(x+2y) - f(x-2y) = 2(f(x+y) - f(x-y)) + 2f(3y) - 6f(2y) + 6f(y). \quad (1.6)$$

It is easy to see that the function $f(x) = ax^3 + bx^2 + cx$ is a solution of the functional equation (1.6). In the present paper we investigate the general solution and the generalized Hyers-Ulam-Rassias stability of the functional equation (1.6).

2. General Solution

In this section we establish the general solution of functional equation (1.6).

Theorem 2.1. Let X, Y be vector spaces, and let $f : X \rightarrow Y$ be a function. Then f satisfies (1.6) if and only if there exists a unique additive function $A : X \rightarrow Y$, a unique symmetric and biadditive function $Q : X \times X \rightarrow Y$, and a unique symmetric and 3-additive function $C : X \times X \times X \rightarrow Y$ such that $f(x) = A(x) + Q(x, x) + C(x, x, x)$ for all $x \in X$.

Proof. Suppose that $f(x) = A(x) + Q(x, x) + C(x, x, x)$ for all $x \in X$, where $A : X \rightarrow Y$ is additive, $Q : X \times X \rightarrow Y$ is symmetric and biadditive, and $C : X \times X \times X \rightarrow Y$ is symmetric and 3-additive. Then it is easy to see that f satisfies (1.6). For the converse let f satisfy (1.6). We decompose f into the even part and odd part by setting

$$f_e(x) = \frac{1}{2}(f(x) + f(-x)), \quad f_o(x) = \frac{1}{2}(f(x) - f(-x)), \quad (2.1)$$

for all $x \in X$. By (1.6), we have

$$\begin{aligned} & f_e(x+2y) - f_e(x-2y) \\ &= \frac{1}{2}[f(x+2y) + f(-x-2y) - f(x-2y) - f(-x+2y)] \\ &= \frac{1}{2}[f(x+2y) - f(x-2y)] + \frac{1}{2}[f((-x) + (-2y)) - f((-x) - (-2y))] \\ &= \frac{1}{2}[2f(x+y) - 2f(x-y) + 2f(3y) - 6f(2y) + 6f(y)] \\ &\quad + \frac{1}{2}[2f(-x-y) - 2f(-x+y) + 2f(-3y) - 6f(-2y) + 6f(-y)] \\ &= 2\left[\frac{1}{2}(f(x+y) + f(-x-y))\right] - 2\left[\frac{1}{2}(f(x-y) + f(-x+y))\right] \\ &\quad + 2\left[\frac{1}{2}(f(3y) + f(-3y))\right] - 6\left[\frac{1}{2}(f(2y) + f(-2y))\right] + 6\left[\frac{1}{2}(f(y) + f(-y))\right] \\ &= 2(f_e(x+y) - f_e(x-y)) + 2f_e(3y) - 6f_e(2y) + 6f_e(y), \end{aligned} \quad (2.2)$$

for all $x, y \in X$. This means that f_e satisfies (1.6), that is,

$$f_e(x+2y) - f_e(x-2y) = 2(f_e(x+y) - f_e(x-y)) + 2f_e(3y) - 6f_e(2y) + 6f_e(y). \quad (2.3)$$

Now putting $x = y = 0$ in (2.3), we get $f_e(0) = 0$. Setting $x = 0$ in (2.3), by evenness of f_e we obtain

$$3f_e(2y) = f_e(3y) + 3f_e(y). \quad (2.4)$$

Replacing x by y in (2.3), we obtain

$$4f_e(2y) = f_e(3y) + 7f_e(y). \quad (2.5)$$

Comparing (2.4) with (2.5), we get

$$f_e(3y) = 9f_e(y). \quad (2.6)$$

By utilizing (2.5) with (2.6), we obtain

$$f_e(2y) = 4f_e(y). \quad (2.7)$$

Hence, according to (2.6) and (2.7), (2.3) can be written as

$$f_e(x+2y) - f_e(x-2y) = 2f_e(x+y) - 2f_e(x-y). \quad (2.8)$$

With the substitution $x := x+y$, $y := x-y$ in (2.8), we have

$$f_e(3x-y) - f_e(x-3y) = 8f_e(x) - 8f_e(y). \quad (2.9)$$

Replacing y by $-y$ in above relation, we obtain

$$f_e(3x+y) - f_e(x+3y) = 8f_e(x) - 8f_e(y). \quad (2.10)$$

Setting $x+y$ instead of x in (2.8), we get

$$f_e(x+3y) - f_e(x-y) = 2f_e(x+2y) - 2f_e(x). \quad (2.11)$$

Interchanging x and y in (2.11), we get

$$f_e(3x+y) - f_e(x-y) = 2f_e(2x+y) - 2f_e(y). \quad (2.12)$$

If we subtract (2.12) from (2.11) and use (2.10), we obtain

$$f_e(x+2y) - f_e(2x+y) = 3f_e(y) - 3f_e(x), \quad (2.13)$$

which, by putting $y := 2y$ and using (2.7), leads to

$$f_e(x+4y) - 4f_e(x+y) = 12f_e(y) - 3f_e(x). \quad (2.14)$$

Let us interchange x and y in (2.14). Then we see that

$$f_e(4x+y) - 4f_e(x+y) = 12f_e(x) - 3f_e(y), \quad (2.15)$$

and by adding (2.14) and (2.15), we arrive at

$$f_e(x+4y) + f_e(4x+y) = 8f_e(x+y) + 9f_e(x) + 9f_e(y). \quad (2.16)$$

Replacing y by $x + y$ in (2.8), we obtain

$$f_e(3x + 2y) - f_e(x + 2y) = 2f_e(2x + y) - 2f_e(y). \quad (2.17)$$

Let us Interchange x and y in (2.17). Then we see that

$$f_e(2x + 3y) - f_e(2x + y) = 2f_e(x + 2y) - 2f_e(x). \quad (2.18)$$

Thus by adding (2.17) and (2.18), we have

$$f_e(2x + 3y) + f_e(3x + 2y) = 3f_e(x + 2y) + 3f_e(2x + y) - 2f_e(x) - 2f_e(y). \quad (2.19)$$

Replacing x by $2x$ in (2.11) and using (2.7) we have

$$f_e(2x + 3y) - f_e(2x - y) = 8f_e(x + y) - 8f_e(x), \quad (2.20)$$

and interchanging x and y in (2.20) yields

$$f_e(3x + 2y) - f_e(x - 2y) = 8f_e(x + y) - 8f_e(y). \quad (2.21)$$

If we add (2.20) to (2.21), we have

$$f_e(2x + 3y) + f_e(3x + 2y) = f_e(2x - y) + f_e(x - 2y) + 16f_e(x + y) - 8f_e(x) - 8f_e(y). \quad (2.22)$$

Interchanging x and y in (2.8), we get

$$f_e(2x + y) - f_e(2x - y) = 2f_e(x + y) - 2f_e(x - y), \quad (2.23)$$

and by adding the last equation and (2.8) with (2.19), we get

$$\begin{aligned} & f_e(2x + 3y) + f_e(3x + 2y) - f_e(2x - y) - f_e(x - 2y) \\ &= 2f_e(x + 2y) + 2f_e(2x + y) + 4f_e(x + y) - 4f_e(x - y) - 2f_e(x) - 2f_e(y). \end{aligned} \quad (2.24)$$

Now according to (2.22) and (2.24), it follows that

$$f_e(x + 2y) + f_e(2x + y) = 6f_e(x + y) + 2f_e(x - y) - 3f_e(x) - 3f_e(y). \quad (2.25)$$

From the substitution $y = -y$ in (2.25) it follows that

$$f_e(x - 2y) + f_e(2x - y) = 6f_e(x - y) + 2f_e(x + y) - 3f_e(x) - 3f_e(y). \quad (2.26)$$

Replacing y by $2y$ in (2.25) we have

$$f_e(x+4y) + 4f_e(x+y) = 6f_e(x+2y) + 2f_e(x-2y) - 3f_e(x) - 12f_e(y), \quad (2.27)$$

and interchanging x and y yields

$$f_e(4x+y) + 4f_e(x+y) = 6f_e(2x+y) + 2f_e(2x-y) - 12f_e(x) - 3f_e(y). \quad (2.28)$$

By adding (2.27) and (2.28) and then using (2.25) and (2.26), we lead to

$$f_e(x+4y) + f_e(4x+y) = 32f_e(x+y) + 24f_e(x-y) - 39f_e(x) - 39f_e(y). \quad (2.29)$$

If we compare (2.16) and (2.29), we conclude that

$$f_e(x+y) + f_e(x-y) = 2f_e(x) + 2f_e(y). \quad (2.30)$$

This means that f_e is quadratic. Thus there exists a unique quadratic function $Q : X \times X \rightarrow Y$ such that $f_e(x) = Q(x, x)$, for all $x \in X$. On the other hand we can show that f_o satisfies (1.6), that is,

$$f_o(x+2y) - f_o(x-2y) = 2(f_o(x+y) - f_o(x-y)) + 2f_o(3y) - 6f_o(2y) + 6f_o(y). \quad (2.31)$$

Now we show that the mapping $g : X \rightarrow Y$ defined by $g(x) := f_o(2x) - 8f_o(x)$ is additive and the mapping $h : X \rightarrow Y$ defined by $h(x) := f_o(2x) - 2f_o(x)$ is cubic. Putting $x = 0$ in (2.31), then by oddness of f_o , we have

$$4f_o(2y) = 5f_o(y) + f_o(3y). \quad (2.32)$$

Hence (2.31) can be written as

$$f_o(x+2y) - f_o(x-2y) = 2f_o(x+y) - 2f_o(x-y) + 2f_o(2y) - 4f_o(y). \quad (2.33)$$

From the substitution $y := -y$ in (2.33) it follows that

$$f_o(x-2y) - f_o(x+2y) = 2f_o(x-y) - 2f_o(x+y) - 2f_o(2y) + 4f_o(y). \quad (2.34)$$

Interchange x and y in (2.33), and it follows that

$$f_o(2x+y) + f_o(2x-y) = 2f_o(x+y) + 2f_o(x-y) + 2f_o(2x) - 4f_o(x). \quad (2.35)$$

With the substitutions $x := x-y$ and $y := x+y$ in (2.35), we have

$$f_o(3x-y) + f_o(x-3y) = 2f_o(2x-2y) - 4f_o(x-y) + 2f_o(2x) - 2f_o(2y). \quad (2.36)$$

Replace x by $x - y$ in (2.34). Then we have

$$f_o(x - 3y) - f_o(x + y) = 2f_o(x - 2y) - 2f_o(x) - 2f_o(2y) + 4f_o(y). \quad (2.37)$$

Replacing y by $-y$ in (2.37) gives

$$f_o(x + 3y) - f_o(x - y) = 2f_o(x + 2y) - 2f_o(x) + 2f_o(2y) - 4f_o(y). \quad (2.38)$$

Interchanging x and y in (2.38), we get

$$f_o(3x + y) + f_o(x - y) = 2f_o(2x + y) - 2f_o(y) + 2f_o(2x) - 4f_o(x). \quad (2.39)$$

If we add (2.38) to (2.39), we have

$$\begin{aligned} f_o(x + 3y) + f_o(3x + y) \\ = 2f_o(x + 2y) + 2f_o(2x + y) + 2f_o(2x) + 2f_o(2y) - 6f_o(x) - 6f_o(y). \end{aligned} \quad (2.40)$$

Replacing y by $-y$ in (2.36) gives

$$f_o(x + 3y) + f_o(3x + y) = 2f_o(2x + 2y) - 4f_o(x + y) + 2f_o(2x) + 2f_o(2y). \quad (2.41)$$

By comparing (2.40) with (2.41), we arrive at

$$f_o(x + 2y) + f_o(2x + y) = f_o(2x + 2y) - 2f_o(x + y) + 3f_o(x) + 3f_o(y). \quad (2.42)$$

Replacing y by $-y$ in (2.42) gives

$$f_o(x - 2y) + f_o(2x - y) = f_o(2x - 2y) - 2f_o(x - y) + 3f_o(x) - 3f_o(y). \quad (2.43)$$

With the substitution $y := x + y$ in (2.43), we have

$$f_o(x - y) - f_o(x + 2y) = -f_o(2y) - 3f_o(x + y) + 3f_o(x) + 2f_o(y), \quad (2.44)$$

and replacing $-y$ by y gives

$$f_o(x + y) - f_o(x - 2y) = f_o(2y) - 3f_o(x - y) + 3f_o(x) - 2f_o(y). \quad (2.45)$$

Let us interchange x and y in (2.45). Then we see that

$$f_o(x + y) + f_o(2x - y) = f_o(2x) + 3f_o(x - y) - 2f_o(x) + 3f_o(y). \quad (2.46)$$

If we add (2.45) to (2.46), we have

$$f_o(2x - y) - f_o(x - 2y) = f_o(2x) - 2f_o(x + y) + f_o(x) + f_o(2y) + f_o(y). \quad (2.47)$$

Adding (2.42) to (2.47) and using (2.33) and (2.35), we obtain

$$f_o(2(x + y)) - 8f_o(x + y) = [f_o(2x) - 8f_o(x)] + [f_o(2y) - 8f_o(y)], \quad (2.48)$$

for all $x, y \in X$. The last equality means that

$$g(x + y) = g(x) + g(y), \quad (2.49)$$

for all $x, y \in X$. Therefore the mapping $g : X \rightarrow Y$ is additive. With the substitutions $x := 2x$ and $y := 2y$ in (2.35), we have

$$f_o(4x + 2y) + f_o(4x - 2y) = 2f_o(2x + 2y) + 2f_o(2x - 2y) + 2f_o(4x) - 4f_o(2x). \quad (2.50)$$

Let $g : X \rightarrow Y$ be the additive mapping defined above. It is easy to show that f_o is cubic-additive function. Then there exists a unique function $C : X \times X \times X \rightarrow Y$ and a unique additive function $A : X \rightarrow Y$ such that $f_o(x) = C(x, x, x) + A(x)$, for all $x \in X$, and C is symmetric and 3-additive. Thus for all $x \in X$, we have

$$f(x) = f_e(x) + f_o(x) = Q(x, x) + C(x, x, x) + A(x). \quad (2.51)$$

This completes the proof of theorem. □

The following corollary is an alternative result of Theorem 2.1.

Corollary 2.2. *Let X, Y be vector spaces, and let $f : X \rightarrow Y$ be a function satisfying (1.6). Then the following assertions hold.*

- (a) *If f is even function, then f is quadratic.*
- (b) *If f is odd function, then f is cubic-additive.*

3. Stability

We now investigate the generalized Hyers-Ulam-Rassias stability problem for functional equation (1.6). From now on, let X be a real vector space, and let Y be a Banach space. Now before taking up the main subject, given $f : X \rightarrow Y$, we define the difference operator $D_f : X \times X \rightarrow Y$ by

$$D_f(x, y) = f(x + 2y) - f(x - 2y) - 2[f(x + y) - f(x - y)] - 2f(3y) + 6f(2y) - 6f(y), \quad (3.1)$$

for all $x, y \in X$. We consider the following functional inequality:

$$\|D_f(x, y)\| \leq \phi(x, y), \tag{3.2}$$

for an upper bound $\phi : X \times X \rightarrow [0, \infty)$.

Theorem 3.1. *Let $s \in \{1, -1\}$ be fixed. Suppose that an even mapping $f : X \rightarrow Y$ satisfies $f(0) = 0$, and*

$$\|D_f(x, y)\| \leq \phi(x, y), \tag{3.3}$$

for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that

$$\sum_{i=0}^{\infty} 4^{si} \left[\phi(2^{-si}x, 2^{-si}x) + \frac{1}{2}\phi(0, 2^{-si}x) \right] < \infty \tag{3.4}$$

and that

$$\lim_n 4^{sn} \phi(2^{-sn}x, 2^{-sn}y) = 0, \tag{3.5}$$

for all $x, y \in X$, then the limit

$$Q(x) := \lim_n 4^{sn} f(2^{-sn}x) \tag{3.6}$$

exists for all $x \in X$, and $Q : X \rightarrow Y$ is a unique quadratic function satisfying (1.6), and

$$\|f(x) - Q(x)\| \leq \frac{1}{8} \sum_{i=(s+1)/2}^{\infty} 4^{si} \left(\phi(2^{-si}x, 2^{-si}x) + \frac{1}{2}\phi(0, 2^{-si}x) \right), \tag{3.7}$$

for all $x \in X$.

Proof. Let $s = 1$. Putting $x = 0$ in (3.3), we get

$$\|2[f(3y) - 3f(2y) + 3f(y)]\| \leq \phi(0, y), \tag{3.8}$$

for all $y \in X$. On the other hand by replacing y by x in (3.3), it follows that

$$\|-f(3y) + 4f(2y) - 7f(y)\| \leq \phi(y, y), \tag{3.9}$$

for all $y \in X$. Combining (3.8) and (3.9), we lead to

$$\|2f(2y) - 8f(y)\| \leq 2\phi(y, y) + \phi(0, y), \tag{3.10}$$

for all $y \in X$. With the substitution $y := x/2$ in (3.10) and then dividing both sides of inequality by 2, we get

$$\left\| f(x) - 4f\left(\frac{x}{2}\right) \right\| \leq \frac{1}{2} \left[2\phi\left(\frac{x}{2}, \frac{x}{2}\right) + \phi\left(0, \frac{x}{2}\right) \right]. \quad (3.11)$$

Now, using methods similar as in [8, 34, 35], we can easily show that the function $Q : X \rightarrow Y$ defined by $Q(x) = \lim_{n \rightarrow \infty} 4^n f(x/2^n)$ for all $x \in X$ is unique quadratic function satisfying (1.6) and (3.7). Let $s = -1$. Then by (3.10) we have

$$\left\| \frac{f(2x)}{4} - f(x) \right\| \leq \frac{1}{8} (2\phi(x, x) + \phi(0, x)), \quad (3.12)$$

for all $x \in X$. And analogously, as in the case $s = 1$, we can show that the function $Q : X \rightarrow Y$ defined by $Q(x) := \lim_{n \rightarrow \infty} 4^{-n} f(2^n x)$ is unique quadratic function satisfying (1.6) and (3.7). \square

Theorem 3.2. *Let $s \in \{1, -1\}$ be fixed. Let $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that*

$$\sum_{i=1}^{\infty} 2^{si} \left[\phi\left(\frac{x}{2^{si}}, \frac{x}{2^{si+1}}\right) + \phi\left(0, \frac{x}{2^{si+1}}\right) \right] < \infty \quad (3.13)$$

and that

$$\lim_{n \rightarrow \infty} 2^{sn} \phi\left(\frac{x}{2^{sn}}, \frac{y}{2^{sn}}\right) = 0, \quad (3.14)$$

for all $x, y \in X$.

Suppose that an odd mapping $f : X \rightarrow Y$ satisfies

$$\|D_f(x, y)\| \leq \phi(x, y), \quad (3.15)$$

for all $x, y \in X$.

Then the limit

$$A(x) := \lim_{n \rightarrow \infty} 2^{sn} \left[f\left(\frac{x}{2^{sn-1}}\right) - 8f\left(\frac{x}{2^{sn}}\right) \right] \quad (3.16)$$

exists, for all $x \in X$, and $A : X \rightarrow Y$ is a unique additive function satisfying (1.6), and

$$\|f(2x) - 8f(x) - A(x)\| \leq \sum_{i=|s-1|/2}^{\infty} 2^{si} \phi\left(\frac{x}{2^{si}}, \frac{x}{2^{si+1}}\right) + 2 \sum_{i=|s-1|/2}^{\infty} 2^{si} \phi\left(0, \frac{x}{2^{si+1}}\right), \quad (3.17)$$

for all $x \in X$.

Proof. Let $s = 1$. set $x = 0$ in (3.15). Then by oddness of f we have

$$\|2f(3y) - 8f(2y) + 16f(y)\| \leq \phi(0, y), \tag{3.18}$$

for all $y \in X$. Replacing x by $2y$ in (3.15) we get

$$\|f(4y) - 4f(3y) + 6f(2y) - 4f(y)\| \leq \phi(2y, y). \tag{3.19}$$

Combining (3.18) and (3.19), we lead to

$$\|f(4y) - 10f(2y) + 16f(y)\| \leq \phi(2y, y) + 2\phi(0, y), \tag{3.20}$$

for all $y \in X$. Putting $y := x/2$ and $g(x) := f(2x) - 8f(x)$, for all $x \in X$. Then we get

$$\|g(x) - 2g\left(\frac{x}{2}\right)\| \leq \phi\left(x, \frac{x}{2}\right) + 2\phi\left(0, \frac{x}{2}\right), \tag{3.21}$$

for all $x \in X$. Now, in a similar way as in [8, 34, 35], we can show that the limit $A(x) := \lim_{n \rightarrow \infty} 2^n g(x/2^n)$ exists, for all $x \in X$, and A is the unique function satisfying (1.6) and (3.17). If $s = -1$, then the proof is analogous. \square

Theorem 3.3. Let $s \in \{1, -1\}$ be fixed. Suppose that an odd mapping $f : X \rightarrow Y$ satisfies

$$\|D_f(x, y)\| \leq \phi(x, y), \tag{3.22}$$

for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that

$$\sum_{i=1}^{\infty} 8^{si} \phi\left(\frac{x}{2^{si}}, \frac{x}{2^{si+1}}\right) + \sum_{i=1}^{\infty} 8^{si} \phi\left(0, \frac{x}{2^{si+1}}\right) < \infty \tag{3.23}$$

and that $\lim_{n \rightarrow \infty} 8^{sn} \phi(x/2^{sn}, y/2^{sn}) = 0$, for all $x, y \in X$, then the limit

$$C(x) := \lim_{n \rightarrow \infty} 8^{sn} \left[f\left(\frac{x}{2^{sn-1}}\right) - 2f\left(\frac{x}{2^{sn}}\right) \right] \tag{3.24}$$

exists, for all $x \in X$, and $C : X \rightarrow Y$ is a unique cubic function satisfying (1.6) and

$$\|f(2x) - 2f(x) - C(x)\| \leq \sum_{i=|s-1|/2}^{\infty} 8^{si} \phi\left(\frac{x}{2^{si}}, \frac{x}{2^{si+1}}\right) + 2 \sum_{i=|s-1|/2}^{\infty} 8^{si} \phi\left(0, \frac{x}{2^{si+1}}\right), \tag{3.25}$$

for all $x \in X$.

Proof. We prove the theorem for $s = 1$. When $s = -1$ we have a similar proof. It is easy to see that f satisfies (3.20). Set $h(x) := f(2x) - 2f(x)$ then by putting $y := x/2$ in (3.20), it follows that

$$\left\| h(x) - 8h\left(\frac{x}{2}\right) \right\| \leq \phi\left(x, \frac{x}{2}\right) + 2\phi\left(0, \frac{x}{2}\right), \quad (3.26)$$

for all $x \in X$. By using (3.26), we may define a mapping $C : X \rightarrow Y$ as $C(x) := \lim_{n \rightarrow \infty} 8^n h(x/2^n)$, for all $x \in X$. Similar to Theorem 3.1, we can show that C is the unique cubic function satisfying (1.6) and (3.25). \square

Theorem 3.4. *Suppose that an odd mapping $f : X \rightarrow Y$ satisfies*

$$\|D_f(x, y)\| \leq \phi(x, y), \quad (3.27)$$

for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that

$$\sum_{i=1}^{\infty} 8^i \phi\left(\frac{x}{2^i}, \frac{x}{2^{i+1}}\right) + \sum_{i=1}^{\infty} 8^i \phi\left(0, \frac{x}{2^{i+1}}\right) < \infty, \quad (3.28)$$

and that $\lim_{n \rightarrow \infty} 8^n \phi(x/2^n, y/2^n) = 0$, for all $x, y \in X$, then there exists a unique cubic function $C : X \rightarrow Y$ and a unique additive function $A : X \rightarrow Y$ such that

$$\|f(x) - C(x) - A(x)\| \leq \frac{1}{6} \sum_{i=0}^{\infty} (2^i + 8^i) \phi\left(\frac{x}{2^i}, \frac{x}{2^{i+1}}\right) + \frac{1}{3} \sum_{i=0}^{\infty} (2^i + 8^i) \phi\left(0, \frac{x}{2^{i+1}}\right), \quad (3.29)$$

for all $x \in X$.

Proof. By Theorems 3.2 and 3.3, there exist an additive mapping $A_o : X \rightarrow Y$ and a cubic mapping $C_o : X \rightarrow Y$ such that

$$\begin{aligned} \|f(2x) - 8f(x) - A_o(x)\| &\leq \sum_{i=|s-1|/2}^{\infty} 2^{si} \phi\left(\frac{x}{2^{si}}, \frac{x}{2^{si+1}}\right) + 2 \sum_{i=|s-1|/2}^{\infty} 2^{si} \phi\left(0, \frac{x}{2^{si+1}}\right), \\ \|f(2x) - 2f(x) - C_o(x)\| &\leq \sum_{i=|s-1|/2}^{\infty} 8^{si} \phi\left(\frac{x}{2^{si}}, \frac{x}{2^{si+1}}\right) + 2 \sum_{i=|s-1|/2}^{\infty} 8^{si} \phi\left(0, \frac{x}{2^{si+1}}\right), \end{aligned} \quad (3.30)$$

for all $x \in X$. Combine the two equations of (3.30) to obtain

$$\left\| f(x) - \frac{1}{6}C_o(x) + \frac{1}{6}A_o(x) \right\| \leq \frac{1}{6} \sum_{i=0}^{\infty} (2^i + 8^i) \phi\left(\frac{x}{2^i}, \frac{x}{2^{i+1}}\right) + \frac{1}{3} \sum_{i=0}^{\infty} (2^i + 8^i) \phi\left(0, \frac{x}{2^{i+1}}\right), \quad (3.31)$$

for all $x \in X$. So we get (3.29) by letting $A(x) = -(1/6)A_0(x)$, and $C(x) = (1/6)C_0(x)$, for all $x \in X$. To prove the uniqueness of A and C , let $A_1, C_1 : X \rightarrow Y$ be another additive and cubic maps satisfying (3.29). Let $A' = A - A_1$, and let $C' = C - C_1$. So

$$\begin{aligned} \|A'(x) - C'(x)\| &\leq \|f(x) - A(x) - C(x)\| + \|f(x) - A_1(x) - C_1(x)\| \\ &\leq 2 \left[\frac{1}{30} \sum_{i=0}^{\infty} (2^i + 8^i) \phi \left(\frac{x}{2^i}, \frac{x}{2^{i+1}} \right) + \frac{1}{15} \sum_{i=0}^{\infty} (2^i + 8^i) \phi \left(0, \frac{x}{2^{i+1}} \right) \right], \end{aligned} \tag{3.32}$$

for all $x \in X$. Since

$$\lim_{n \rightarrow \infty} \left\{ \sum_{i=1}^{\infty} 8^{i+n} \phi \left(\frac{x}{2^{i+n}}, \frac{x}{2^{i+n+1}} \right) + \sum_{i=1}^{\infty} 8^{i+n} \phi \left(0, \frac{x}{2^{i+n+1}} \right) \right\} = 0, \tag{3.33}$$

then

$$\lim_{n \rightarrow \infty} \left\{ \sum_{i=1}^{\infty} 2^{i+n} \phi \left(\frac{x}{2^{i+n}}, \frac{x}{2^{i+n+1}} \right) + \sum_{i=1}^{\infty} 2^{i+n} \phi \left(0, \frac{x}{2^{i+n+1}} \right) \right\} = 0, \tag{3.34}$$

for all $x \in X$. Hence (3.32) implies that

$$\lim_{n \rightarrow \infty} 8^n \left\| A' \left(\frac{x}{2^n} \right) - C' \left(\frac{x}{2^n} \right) \right\| = 0, \tag{3.35}$$

for all $x \in X$. On the other hand C and C_1 are cubic, then $C'(x/2^n) = (1/8^n)C'(x)$. Therefore by (3.35) we obtain that $A'(x) = 0$, for all $x \in X$. Again by (3.35) we have $C'(x) = 0$, for all $x \in X$. \square

Theorem 3.5. *Suppose that an odd mapping $f : X \rightarrow Y$ satisfies*

$$\|D_f(x, y)\| \leq \phi(x, y), \tag{3.36}$$

for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that

$$\sum_{i=1}^{\infty} \frac{1}{2^i} \phi(2^i x, 2^{i-1} x) + \sum_{i=1}^{\infty} 2^i \phi(0, 2^{i-1} x) < \infty \tag{3.37}$$

and that $\lim_{n \rightarrow \infty} (1/2^n)\phi(2^n x, 2^n y) = 0$, for all $x, y \in X$, then there exist a unique cubic function $C : X \rightarrow Y$ and a unique additive function $A : X \rightarrow Y$ such that

$$\begin{aligned} &\|f(x) - C(x) - A(x)\| \\ &\leq \frac{1}{30} \sum_{i=1}^{\infty} \left(\frac{1}{2^i} + \frac{1}{8^i} \right) \left(\phi(2^i x, 2^{i-1} x) \right) + \frac{1}{15} \sum_{i=1}^{\infty} \left(\frac{1}{2^i} + \frac{1}{8^i} \right) \left(\phi(0, 2^{i-1} x) \right), \end{aligned} \tag{3.38}$$

for all $x \in X$.

Proof. The proof is similar to the proof of Theorem 3.4. \square

Now we establish the generalized Hyers-Ulam-Rassias stability of functional equation (1.6) as follows.

Theorem 3.6. *Suppose that a mapping $f : X \rightarrow Y$ satisfies $f(0) = 0$ and $\|D_f(x, y)\| \leq \phi(x, y)$, for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that*

$$\sum_{i=0}^{\infty} \left\{ 8^i \left[\phi\left(\frac{x}{2^i}, \frac{x}{2^{i+1}}\right) + \phi\left(0, \frac{x}{2^{i+1}}\right) \right] + 4^i \phi\left(\frac{x}{2^i}, \frac{x}{2^i}\right) \right\} < \infty \quad (3.39)$$

and that $\lim_{n \rightarrow \infty} 8^n \phi(x/2^n, y/2^n) = 0$, for all $x, y \in X$, then there exist a unique additive function $A : X \rightarrow Y$ a unique quadratic function $Q : X \rightarrow Y$ and a unique cubic function $C : X \rightarrow Y$ such that

$$\begin{aligned} & \|f(x) - A(x) - Q(x) - C(x)\| \\ & \leq \frac{1}{6} \sum_{i=0}^{\infty} (2^i + 8^i) \left[\phi\left(\frac{x}{2^i}, \frac{x}{2^{i+1}}\right) + 2\phi\left(0, \frac{x}{2^{i+1}}\right) \right] + \frac{1}{8} \sum_{i=1}^{\infty} 4^i \left[\phi\left(\frac{x}{2^i}, \frac{x}{2^i}\right) + \frac{1}{2}\phi\left(0, \frac{x}{2^i}\right) \right], \end{aligned} \quad (3.40)$$

for all $x \in X$.

Proof. Let $f_e(x) = (1/2)(f(x) + f(-x))$, for all $x \in X$. Then $f_e(0) = 0$, $f_e(-x) = f_e(x)$, and $\|D_{f_e}(x, y)\| \leq (1/2)[\phi(x, y) + \phi(-x, -y)]$, for all $x, y \in X$. Hence in view of Theorem 3.1 there exists a unique quadratic function $Q : X \rightarrow Y$ satisfying (3.7). Let $f_o(x) = (1/2)(f(x) - f(-x))$, for all $x \in X$. Then $f_o(0) = 0$, $f_o(-x) = -f_o(x)$, and $\|D_{f_o}(x, y)\| \leq (1/2)[\phi(x, y) + \phi(-x, -y)]$, for all $x, y \in X$. From Theorem 3.4, it follows that there exist a unique cubic function $C : X \rightarrow Y$ and a unique additive function $A : X \rightarrow Y$ satisfying (3.29). Now it is obvious that (3.40) holds true for all $x \in X$, and the proof of theorem is complete. \square

Corollary 3.7. *Let $p + q > 3$, $\theta \geq 0$. Suppose that a mapping $f : X \rightarrow Y$ satisfies $f(0) = 0$, and*

$$\|D_f(x, y)\| \leq \theta(\|x\|^p \|y\|^q), \quad (3.41)$$

for all $x, y \in X$. Then there exist a unique additive function $A : X \rightarrow Y$, a unique quadratic function $Q : X \rightarrow Y$, and a unique cubic function $C : X \rightarrow Y$ satisfying

$$\|f(x) - A(x) - Q(x) - C(x)\| \leq \theta \|x\|^{p+q} \left[\left(\frac{1}{6 \times 2^q} \right) \left(\frac{2}{2 - 2^{p+q}} + \frac{8}{8 - 2^{p+q}} \right) + \frac{1}{8} \left(\frac{2^{p+q}}{4 - 2^{p+q}} \right) \right], \quad (3.42)$$

for all $x \in X$.

Proof. It follows from Theorem 3.6 by taking $\phi(x, y) = \theta(\|x\|^p \|y\|^q)$, for all $x, y \in X$. \square

Theorem 3.8. *Suppose that $f : X \rightarrow Y$ satisfies $f(0) = 0$, and $\|D_f(x, y)\| \leq \phi(x, y)$, for all $x, y \in X$. If the upper bound $\phi : X \times X \rightarrow [0, \infty)$ is a mapping such that*

$$\sum_{i=1}^{\infty} \left\{ \frac{1}{2^i} [\phi(2^i x, 2^{i-1} x) + \phi(0, 2^{i-1} x)] + \frac{1}{4^i} \phi(2^i x, 2^i x) \right\} < \infty \tag{3.43}$$

and that $\lim_{n \rightarrow \infty} (1/2^n)\phi(2^n x, 2^n y) = 0$, for all $x, y \in X$, then there exists a unique additive function $A : X \rightarrow Y$, a unique quadratic function $Q : X \rightarrow Y$, and a unique cubic function $C : X \rightarrow Y$ such that

$$\begin{aligned} & \|f(x) - A(x) - Q(x) - C(x)\| \\ & \leq \frac{1}{6} \left[\sum_{i=1}^{\infty} \left(\frac{1}{2^i} + \frac{1}{8^i} \right) (\phi(2^i x, 2^{i-1} x) + 2\phi(0, 2^{i-1} x)) \right] + \frac{1}{8} \sum_{i=0}^{\infty} \frac{1}{4^i} \left[\phi(2^i x, 2^i x) + \frac{1}{2} \phi(0, 2^i x) \right], \end{aligned} \tag{3.44}$$

for all $x \in X$.

By Theorem 3.8, we are going to investigate the following stability problem for functional equation (1.6).

Corollary 3.9. *Let $p + q < 1$, $\theta > 0$. Suppose that $f : X \rightarrow Y$ satisfies $f(0) = 0$, and*

$$\|D_f(x, y)\| \leq \theta(\|x\|^p \|y\|^q), \tag{3.45}$$

for all $x, y \in X$, then there exist a unique additive function $A : X \rightarrow Y$, a unique quadratic function $Q : X \rightarrow Y$, and a unique cubic function $C : X \rightarrow Y$ satisfying

$$\begin{aligned} & \|f(x) - A(x) - Q(x) - C(x)\| \\ & \leq \theta \|x\|^{p+q} \left\{ \left(\frac{1}{6 \times 2^q} \right) \left(\frac{2^{p+q}}{2 - 2^{p+q}} + \frac{2^{p+q}}{8 - 2^{p+q}} \right) + \frac{1}{8 - 2^{p+q+3}} \right\}, \end{aligned} \tag{3.46}$$

for all $x \in X$.

By Corollary 3.9, we solve the following Hyers-Ulam stability problem for functional equation (1.6).

Corollary 3.10. *Let ϵ be a positive real number. Suppose that a mapping $f : X \rightarrow Y$ satisfies $f(0) = 0$, and $\|D_f(x, y)\| \leq \epsilon$, for all $x, y \in X$, then there exist a unique additive function $A : X \rightarrow Y$, a unique quadratic function $Q : X \rightarrow Y$, and a unique cubic function $C : X \rightarrow Y$ such that*

$$\|f(x) - A(x) - Q(x) - C(x)\| \leq \frac{5}{14} \epsilon, \tag{3.47}$$

for all $x \in X$.

Acknowledgments

The authors would like to express their sincere thanks to referees for their invaluable comments. The first author would like to thank the Semnan University for its financial support. Also, the fourth author would like to thank the office of gifted students at Semnan University for its financial support.

References

- [1] S. M. Ulam, *Problems in Modern Mathematics*, chapter 6, John Wiley & Sons, New York, NY, USA, 1940.
- [2] D. H. Hyers, G. Isac, and Th. M. Rassias, *Stability of Functional Equations in Several Variables*, Progress in Nonlinear Differential Equations and Their Applications, 34, Birkhäuser, Boston, Mass, USA, 1998.
- [3] D. G. Bourgin, "Classes of transformations and bordering transformations," *Bulletin of the American Mathematical Society*, vol. 57, pp. 223–237, 1951.
- [4] T. Aoki, "On the stability of the linear transformation in Banach spaces," *Journal of the Mathematical Society of Japan*, vol. 2, pp. 64–66, 1950.
- [5] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, no. 2, pp. 297–300, 1978.
- [6] J. Aczél and J. Dhombres, *Functional Equations in Several Variables*, vol. 31 of *Encyclopedia of Mathematics and Its Applications*, Cambridge University Press, Cambridge, UK, 1989.
- [7] P. W. Cholewa, "Remarks on the stability of functional equations," *Aequationes Mathematicae*, vol. 27, no. 1-2, pp. 76–86, 1984.
- [8] G.-L. Forti, "Comments on the core of the direct method for proving Hyers-Ulam stability of functional equations," *Journal of Mathematical Analysis and Applications*, vol. 295, no. 1, pp. 127–133, 2004.
- [9] Z. Gajda, "On stability of additive mappings," *International Journal of Mathematics and Mathematical Sciences*, vol. 14, no. 3, pp. 431–434, 1991.
- [10] P. Găvruta, "A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings," *Journal of Mathematical Analysis and Applications*, vol. 184, no. 3, pp. 431–436, 1994.
- [11] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [12] G. Isac and Th. M. Rassias, "On the Hyers-Ulam stability of ψ -additive mappings," *Journal of Approximation Theory*, vol. 72, no. 2, pp. 131–137, 1993.
- [13] K.-W. Jun and H.-M. Kim, "The generalized Hyers-Ulam-Rassias stability of a cubic functional equation," *Journal of Mathematical Analysis and Applications*, vol. 274, no. 2, pp. 267–278, 2002.
- [14] L. Maligranda, "A result of Tosio Aoki about a generalization of Hyers-Ulam stability of additive functions—a question of priority," *Aequationes Mathematicae*, vol. 75, no. 3, pp. 289–296, 2008.
- [15] J. M. Rassias, "On a new approximation of approximately linear mappings by linear mappings," *Discussiones Mathematicae*, vol. 7, pp. 193–196, 1985.
- [16] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Bulletin des Sciences Mathématiques. Deuxième Série*, vol. 108, no. 4, pp. 445–446, 1984.
- [17] J. M. Rassias, "Complete solution of the multi-dimensional problem of Ulam," *Discussiones Mathematicae*, vol. 14, pp. 101–107, 1994.
- [18] J. M. Rassias, "On the stability of a multi-dimensional Cauchy type functional equation," in *Geometry, Analysis and Mechanics*, pp. 365–376, World Scientific, River Edge, NJ, USA, 1994.
- [19] J. M. Rassias, "Solution of a stability problem of Ulam," in *Functional Analysis, Approximation Theory and Numerical Analysis*, pp. 241–249, World Scientific, River Edge, NJ, USA, 1994.
- [20] J. M. Rassias, "Solution of a stability problem of Ulam," *Discussiones Mathematicae*, vol. 12, pp. 95–103, 1992.
- [21] J. M. Rassias, "Solution of a problem of Ulam," *Journal of Approximation Theory*, vol. 57, no. 3, pp. 268–273, 1989.
- [22] J. M. Rassias, "On approximation of approximately linear mappings by linear mappings," *Journal of Functional Analysis*, vol. 46, no. 1, pp. 126–130, 1982.
- [23] Th. M. Rassias, Ed., *Functional Equations and Inequalities*, vol. 518 of *Mathematics and Its Applications*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2000.

- [24] Th. M. Rassias, "On the stability of functional equations in Banach spaces," *Journal of Mathematical Analysis and Applications*, vol. 251, no. 1, pp. 264–284, 2000.
- [25] A. Grabiec, "The generalized Hyers-Ulam stability of a class of functional equations," *Publicationes Mathematicae Debrecen*, vol. 48, no. 3-4, pp. 217–235, 1996.
- [26] J. M. Rassias and M. J. Rassias, "Refined Ulam stability for Euler-Lagrange type mappings in Hilbert spaces," *International Journal of Applied Mathematics & Statistics*, vol. 7, pp. 126–132, 2007.
- [27] J. M. Rassias, "On the stability of the Euler-Lagrange functional equation," *Chinese Journal of Mathematics*, vol. 20, no. 2, pp. 185–190, 1992.
- [28] F. Skof, "Proprieta' locali e approssimazione di operatori," *Rendiconti del Seminario Matematico e Fisico di Milano*, vol. 53, pp. 113–129, 1983.
- [29] J. M. Rassias, "Solution of the Ulam stability problem for cubic mappings," *Glasnik Matematički. Serija III*, vol. 36(56), no. 1, pp. 63–72, 2001.
- [30] J. M. Rassias, "Solution of the Ulam problem for cubic mappings," *Analele Universității din Timișoara. Seria Matematică-Informatică*, vol. 38, no. 1, pp. 121–132, 2000.
- [31] M. E. Gordji, "Stability of a functional equation deriving from quartic and additive functions," to appear in *Bulletin of the Korean Mathematical Society*.
- [32] M. E. Gordji, A. Ebadian, and S. Zolfaghari, "Stability of a functional equation deriving from cubic and quartic functions," *Abstract and Applied Analysis*, vol. 2008, Article ID 801904, 17 pages, 2008.
- [33] M. E. Gordji and H. Khodaei, "Solution and stability of generalized mixed type cubic, quadratic and additive functional equation in quasi-Banach spaces," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 71, pp. 5629–5643, 2009.
- [34] J. Brzdęk and A. Pietrzyk, "A note on stability of the general linear equation," *Aequationes Mathematicae*, vol. 75, no. 3, pp. 267–270, 2008.
- [35] A. Pietrzyk, "Stability of the Euler-Lagrange-Rassias functional equation," *Demonstratio Mathematica*, vol. 39, no. 3, pp. 523–530, 2006.