

*Research Article*

# Exponential Stability and Global Attractors for a Thermoelastic Bresse System

**Zhiyong Ma**

*College of Science, Shanghai Second Polytechnic University, Shanghai 201209, China*

Correspondence should be addressed to Zhiyong Ma, mazhiyong1980@hotmail.com

Received 13 September 2010; Accepted 29 October 2010

Academic Editor: E. Thandapani

Copyright © 2010 Zhiyong Ma. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the stability properties for thermoelastic Bresse system which describes the motion of a linear planar shearable thermoelastic beam. The system consists of three wave equations and two heat equations coupled in certain pattern. The two wave equations about the longitudinal displacement and shear angle displacement are effectively damped by the dissipation from the two heat equations. We use multiplier techniques to prove the exponential stability result when the wave speed of the vertical displacement coincides with the wave speed of the longitudinal or of the shear angle displacement. Moreover, the existence of the global attractor is firstly achieved.

## 1. Introduction

In this paper, we will consider the following system:

$$\rho h w_{1tt} = (Eh(w_{1x} - kw_3) - \alpha\theta_{1t})_x - kGh(\phi_2 + w_{3x} + kw_1), \quad (1.1)$$

$$\rho h w_{3tt} = Gh(\phi_2 + w_{3x} + kw_1)_x + kEh(w_{1x} - kw_3) - k\alpha\theta_{1t}, \quad (1.2)$$

$$\rho I \phi_{2tt} = EI\phi_{2xx} - Gh(\phi_2 + w_{3x} + kw_1) - \alpha\theta_{3x}, \quad (1.3)$$

$$\rho c \theta_{1tt} = \theta_{1xxt} + \theta_{1xx} - \alpha T_0(w_{1tx} - kw_{3t}), \quad (1.4)$$

$$\rho c \theta_{3t} = \theta_{3xx} - \alpha T_0 \phi_{2tx}, \quad (1.5)$$

together with initial conditions

$$\begin{aligned} w_1(x, 0) &= u_0(x), & w_{1t}(x, 0) &= v_0(x), & \phi_2(x, 0) &= \phi_0(x), \\ \phi_{2t}(x, 0) &= \psi_0(x), & w_3(x, 0) &= w_0(x), & w_{3t}(x, 0) &= \varphi_0(x), \\ \theta_1(x, 0) &= \theta_0(x), & \theta_{1t}(x, 0) &= \eta_0(x), & \theta_3(x, 0) &= \xi_0(x) \end{aligned} \quad (1.6)$$

and boundary conditions

$$w_1(x, t) = w_{3x}(x, t) = \phi_2(x, t) = \theta_1(x, t) = \theta_3(x, t) = 0, \quad \text{for } x = 0, 1, \quad (1.7)$$

where  $w_1$ ,  $w_3$ , and  $\phi_2$  are the longitudinal, vertical, and shear angle displacement,  $\theta_1$  and  $\theta_3$  are the temperature deviations from the  $T_0$  along the longitudinal and vertical directions,  $E$ ,  $G$ ,  $\rho$ ,  $I$ ,  $m$ ,  $k$ ,  $h$ , and  $c$  are positive constants for the elastic and thermal material properties.

From this seemingly complicated system, very interesting special cases can be obtained. In particular, the isothermal system is exactly the system obtained by Bresse [1] in 1856. The Bresse system, (1.1)–(1.3) with  $\theta_1$ ,  $\theta_3$  removed, is more general than the well-known Timoshenko system where the longitudinal displacement  $w_1$  is not considered. If both  $\theta_1$  and  $w_1$  are neglected, the Bresse thermoelastic system simplifies to the following Timoshenko thermoelastic system:

$$\begin{aligned} \rho h w_{3tt} &= Gh(\phi_2 + w_{3x})_x, \\ \rho I \phi_{2tt} &= EI \phi_{2xx} - Gh(\phi_2 + w_{3x}) - \alpha \theta_{3tx}, \\ \rho c \theta_{3tt} &= \theta_{3xxt} + \theta_{3xx} - \alpha T_0 \phi_{2tx}, \end{aligned} \quad (1.8)$$

which was studied by Messaoudi and Said-Houari [2]. For the boundary conditions

$$w_3(x, t) = \phi_2(x, t) = \theta_{3x}(x, t) = 0, \quad \text{at } x = 0, l, \quad (1.9)$$

they obtained exponential stability for the thermoelastic Timoshenko system (1.8) when  $E = G$ ; later, they proved energy decay for a Timoshenko-type system with history in thermoelasticity of type III [3], and this paper is similar to [2] with an extra damping that comes from the presence of a history term; it improves the result of [2] in the sense that the case of nonequal wave speed has been considered and the relaxation function  $g$  is allowed to decay exponentially or polynomially. We refer the reader to [4–10] for the Timoshenko system with other kinds of damping mechanisms such as viscous damping, viscoelastic damping of Boltzmann type acting on the motion equation of  $w_3$  or  $\phi_2$ . In all these cases, the rotational displacement  $\phi_2$  of the Timoshenko system is effectively damped due to the thermal energy dissipation. In fact, the energy associated with this component of motion decays exponentially. The transverse displacement  $w_3$  is only indirectly damped through the coupling, which can be observed from (1.2). The effectiveness of this damping depends on the type of coupling and the wave speeds. When the wave speeds are the same ( $E = G$ ), the indirect damping is actually strong enough to induce exponential stability for the Timoshenko system, but when the wave speeds are different, the Timoshenko system loses the exponential stability. This phenomenon has been observed for partially damped second-order evolution equations. We would like to mention other works in [11–15] for other related models.

Recently, Liu and Rao [16] considered a similar system; they used semigroup method and showed that the exponentially decay rate is preserved when the wave speed of the vertical displacement coincides with the wave speed of longitudinal displacement or of the shear angle displacement. Otherwise, only a polynomial-type decay rate can be obtained; their main tools are the frequency-domain characterization of exponential decay obtained by Prüss [17] and Huang [18] and of polynomial decay obtained recently by Muñoz Rivera and Fernández Sare [5]. For the attractors, we refer to [19–24].

In this paper, we consider system (1.1)–(1.5); that is, we use multiplier techniques to prove the exponential stability result only for  $E = G$ . However, from the theory of elasticity,  $E$  and  $G$  denote Young's modulus and the shear modulus, respectively. These two elastic moduli are not equal since

$$G = \frac{E}{2(1 + \nu)}, \quad (1.10)$$

where  $\nu \in (0, 1/2)$  is the Poisson's ratio. Thus, the exponential stability for the case of  $E = G$  is only mathematically sound. However, it does provide useful insight into the study of similar models arising from other applications.

## 2. Equal Wave Speeds Case: $E = G$

Here we state and prove a decay result in the case of equal wave speeds propagation.

Define the state spaces

$$\mathcal{H} = H_0^1 \times H_*^1 \times H_0^1 \times H_0^1 \times (L^2)^5, \quad (2.1)$$

where

$$H_*^1 = \left\{ f \in H^1(0, 1) \mid \int_0^1 f(x) = 0 \right\}. \quad (2.2)$$

The associated energy term is given by

$$\begin{aligned} E(t) = & \frac{1}{2} \int_0^1 \left( [Eh(w_{1x} - kw_3)^2 + Gh(\phi_2 + w_{3x} + kw_1)^2 + EI\phi_{2x}^2] \right. \\ & \left. + [\rho h(w_{1t}^2 + w_{3t}^2) + \rho I\phi_{2t}^2] + \frac{\rho c}{T_0} (\theta_{1t}^2 + \theta_{1x}^2 + \theta_3^2) \right) dx. \end{aligned} \quad (2.3)$$

By a straightforward calculation, we have

$$\frac{dE(t)}{dt} = -\frac{1}{T_0} (\|\theta_{1xt}\|^2 + \|\theta_{3x}\|^2). \quad (2.4)$$

From semigroup theory [25, 26], we have the following existence and regularity result; for the explicit proofs, we refer the reader to [16].

**Lemma 2.1.** Let  $u_0(x), w_0(x), \varphi_0(x), \theta_0(x), v_0(x), \phi_0(x), \psi_0(x), \eta_0(x), \xi_0(x) \in \mathcal{L}$  be given. Then problem (1.1)–(1.5) has a unique global weak solution  $(\varphi, \psi, \theta)$  verifying

$$\begin{aligned} w_3(x, t) &\in C(R^+, H_*^1(0, 1)) \cap C^1(R^+, L^2(0, 1)), \\ (w_1(x, t), \phi_2(x, t), \theta_1(x, t), \theta_3(x, t)) &\in C(R^+, H_0^1(0, 1)) \cap C^1(R^+, L^2(0, 1)). \end{aligned} \quad (2.5)$$

We are now ready to state our main stability result.

**Theorem 2.2.** Suppose that  $E = G$  and  $u_0(x), w_0(x), \varphi_0(x), \theta_0(x), v_0(x), \phi_0(x), \psi_0(x), \eta_0(x), \xi_0(x) \in \mathcal{L}$ . Then the energy  $E(t)$  decays exponentially as time tends to infinity; that is, there exist two positive constants  $C$  and  $\mu$  independent of the initial data and  $t$ , such that

$$E(t) \leq CE(0)e^{-\mu t}, \quad \forall t > 0. \quad (2.6)$$

The proof of our result will be established through several lemmas.  
Let

$$I_1 = \int_0^1 \rho I \phi_{2t} \phi_2 + \rho h w_{3t} f, \quad (2.7)$$

where  $f$  is the solution of

$$-f_{xx} = \phi_{2x}, \quad f(0) = f(1) = 0. \quad (2.8)$$

**Lemma 2.3.** Letting  $w_1, w_3, \phi_2, \theta_1, \theta_3$  be a solution of (1.1)–(1.5), then one has, for all  $\varepsilon_1 > 0$ ,

$$\begin{aligned} \frac{I_1(t)}{dt} &\leq -\frac{EI}{2} \|\phi_{2x}\|^2 + \rho I \|\phi_{2t}\|^2 + \varepsilon_1 \left( \|w_{3t}\|^2 + \|(w_{1x} - kw_3)\|^2 \right) \\ &\quad + C(\varepsilon_1) \left( \|\theta_{3x}\|^2 + \|\theta_{1xt}\|^2 + \|\phi_{2t}\|^2 \right). \end{aligned} \quad (2.9)$$

*Proof.*

$$\begin{aligned} \frac{dI_1}{dt} &= -EI \|\phi_{2x}\|^2 + \rho I \|\phi_{2t}\|^2 - \int_0^1 \alpha \theta_{3x} \phi_2 dx \\ &\quad - kEh \int_0^1 (w_{1x} - kw_3) f dx - k\alpha \int_0^1 \theta_{1t} f dx + \rho h \int_0^1 w_{3t} f_t dx, \end{aligned} \quad (2.10)$$

By using the inequalities

$$\begin{aligned} \int_0^1 f_x^2 dx &\leq \int_0^1 \phi_2^2 dx \leq \int_0^1 \phi_{2x}^2 dx, \\ \int_0^1 f_t^2 dx &\leq \int_0^1 f_{tx}^2 dx \leq \int_0^1 \phi_{2t}^2 dx, \end{aligned} \quad (2.11)$$

and Young's inequality, the assertion of the lemma follows.

Let

$$I_2 = \rho c \rho h \int_0^1 \left( \int_0^x \theta_{1t} dy \right) w_{1t} dx. \tag{2.12}$$

□

**Lemma 2.4.** *Letting  $w_1, w_3, \phi_2, \theta_1, \theta_3$  be a solution of (1.1)–(1.5), then one has, for all  $\varepsilon_2 > 0$ ,*

$$\begin{aligned} \frac{dI_2(t)}{dt} &\leq \frac{-\alpha \rho h T_0}{2} \int_0^1 w_{1t}^2 dx + C(\varepsilon_2) \left( \|\theta_{1xt}\|^2 + \|w_{3t}\|^2 \right) \\ &\quad + \varepsilon_2 \left( \|(w_{1x} - kw_3)\|^2 + \|\phi_2 + w_{3x} + kw_1\|^2 \right). \end{aligned} \tag{2.13}$$

*Proof.* Using (1.4) and (1.1), we get

$$\begin{aligned} \frac{I_2(t)}{dt} &= \rho c \rho h \int_0^1 \left( \int_0^x \theta_{1tt} dy \right) w_{1t} dx + \rho c \rho h \int_0^1 \left( \int_0^x \theta_{1t} dy \right) w_{1tt} dx \\ &= \rho h \int_0^1 \left( \int_0^x \theta_{1xxt} + \theta_{1xx} - \alpha T_0 (w_{1tx} - kw_{3t}) dy \right) w_{1t} dx \\ &\quad + \int_0^1 \left( \int_0^x \theta_{1t} dy \right) \left( (Eh(w_{1x} - kw_3) - \alpha \theta_{1t})_x - KGh(\phi_2 + w_{3x} + kw_1) \right) dx \\ &= \rho h \int_0^1 (\theta_{1xt} + \theta_{1x}) w_{1t} dx - \rho h \alpha T_0 \int_0^1 w_{1t}^2 dx + \rho h k \int_0^1 \left( \int_0^x w_{3t} dy \right) w_{1t} dx \\ &\quad + \rho h E h \int_0^1 \left( \theta_{1xt} w_1 + k \theta_{1t} w_3 + \alpha \theta_{1t}^2 \right) dx \\ &\quad - \rho c k G h \int_0^1 \left( \int_0^x \theta_{1t} dy \right) (\phi_2 + w_{3x} + kw_1) dx. \end{aligned} \tag{2.14}$$

The assertion of the lemma then follows, using Young’s and Poincaré’s inequalities.

Let

$$I_3 = \rho c \rho I \int_0^1 \left( \int_0^x \theta_3 dy \right) \phi_{2t} dx. \tag{2.15}$$

□

**Lemma 2.5.** *Letting  $w_1, w_3, \phi_2, \theta_1, \theta_3$  be a solution of (1.1)–(1.5), then one has, for all  $\varepsilon_3 > 0$ ,*

$$\frac{dI_3}{dt} \leq -\frac{\alpha \rho I T_0}{2} \|\phi_{2t}\|^2 + C(\varepsilon_3) \|\theta_{3x}\|^2 + \varepsilon_3 \|\phi_{2x}\|^2 + \varepsilon_3 \|\phi_2 + w_{3x} + kw_1\|^2. \tag{2.16}$$

*Proof.* Using (1.3) and (1.5), we have

$$\begin{aligned}
 \frac{dI_3}{dt} &= \rho c \rho I \int_0^1 \left( \int_0^x \theta_{3t} dy \right) \phi_{2t} dx + \rho c \rho I \int_0^1 \left( \int_0^x \theta_3 dy \right) \phi_{2t} dx \\
 &= \rho I \int_0^1 \int_0^x (\theta_{3xx} - \alpha T_0 \phi_{2xt}) dy \phi_{2t} dx \\
 &\quad + \rho c \int_0^1 \left( \int_0^x \theta_3 dy \right) (EI \phi_{2xx} - Gh(\phi_2 + w_{3x} + kw_1) - \alpha \theta_{3x}) dx \quad (2.17) \\
 &= \rho I \int_0^1 \theta_{3x} \phi_{2t} dx - \alpha IT_0 \int_0^1 \phi_{2t}^2 dx + \rho c EI \int_0^1 \theta_3 \phi_{2x} dx \\
 &\quad - \rho c Gh \int_0^1 \left( \int_0^x \theta_3 dy \right) (\phi_2 + w_{3x} + kw_1) dx - \alpha \rho c \int_0^1 \theta_3^2 dx.
 \end{aligned}$$

Then, using Young's and Poincaré's inequalities, we can obtain the assertion.

Next, we set

$$I_4 = h \rho I \int_0^1 \phi_{2t} (\phi_2 + w_{3x} + kw_1) dx + h \rho I \int_0^1 \phi_{2x} w_{3t} dx. \quad (2.18)$$

□

**Lemma 2.6.** *Letting  $w_1, w_3, \phi_2, \theta_1, \theta_3$  be a solution of (1.1)–(1.5), then one has, for all  $\varepsilon_4 > 0$ ,*

$$\begin{aligned}
 \frac{dI_4}{dt} &\leq -\frac{Gh^2}{2} \int_0^1 (\phi_2 + w_{3x} + kw_1)^2 dx + C(\varepsilon_4) (\|\theta_{3x}\|^2 + \|\theta_{1xt}\|^2) \\
 &\quad + \frac{kh\rho I}{2} (\|\phi_{2t}\|^2 + \|w_{1t}\|^2) + C(\varepsilon_4) \|\phi_{2x}\|^2 + \varepsilon_4 \|w_{1x} - kw_3\|^2. \quad (2.19)
 \end{aligned}$$

*Proof.* Letting (1) =  $I \int_0^1 \phi_{2t} (\phi_2 + w_{3x} + kw_1) dx$ , (2) =  $h \rho I \int_0^1 \phi_{2x} w_{3t} dx$ , then using (1.2), (1.3), we have

$$\begin{aligned}
 (1)' &= \rho I \int_0^1 \phi_{2t} (\phi_2 + w_{3x} + kw_1) dx + h \rho I \int_0^1 \phi_{2t} (\phi_2 + w_{3x} + kw_1)_t dx \\
 &= hEI \int_0^1 \phi_{2xx} (\phi_2 + w_{3x} + kw_1) dx - Gh^2 \int_0^1 (\phi_2 + w_{3x} + kw_1)^2 dx \\
 &\quad - \alpha h \int_0^1 \theta_{3x} (\phi_2 + w_{3x} + kw_1) dx + h \rho I \int_0^1 \phi_{2t}^2 dx + h \rho I \int_0^1 \phi_{2t} (w_{3x} + kw_1)_t dx,
 \end{aligned}$$

$$\begin{aligned}
 (2)' &= I\rho h \int_0^1 \phi_{2xt} w_{3t} dx + I\rho h \int_0^1 \phi_{2x} w_{3xt} dx \\
 &= -I\rho h \int_0^1 \phi_{2t} w_{3xt} dx + IGh \int_0^1 \phi_{2x} (\phi_2 + w_{3x} + kw_1)_x dx \\
 &\quad + IkEh \int_0^1 \phi_{2x} (w_{1x} - kw_3) dx - \alpha Ik \int_0^1 \phi_{2x} \theta_{1t} dx.
 \end{aligned}
 \tag{2.20}$$

Noticing that  $E = G$ , then

$$\begin{aligned}
 I_4' &= (1)' + (2)' \\
 &= -Gh^2 \int_0^1 (\phi_2 + w_{3x} + kw_1)^2 dx - \alpha h \int_0^1 \theta_{3x} (\phi_2 + w_{3x} + kw_1) dx + h\rho I \int_0^1 \phi_{2t}^2 dx \\
 &\quad + khI\rho \int_0^1 \phi_{2t} w_{1t} dx + IkEh \int_0^1 \phi_{2x} (w_{1x} - kw_3) dx - \alpha Ik \int_0^1 \phi_{2x} \theta_{1t} dx.
 \end{aligned}
 \tag{2.21}$$

Then, using Young's inequality, we can obtain the assertion.

We set

$$I_5 = -h\rho \int_0^1 w_{3t} (w_{1x} - kw_3) dx - h\rho \int_0^1 w_{1t} (\phi_2 + w_{3x} + kw_1) dx.
 \tag{2.22}$$

□

**Lemma 2.7.** *Letting  $w_1, w_3, \phi_2, \theta_1, \theta_3$  be a solution of (1.1)–(1.5), then one has, for all  $\varepsilon_5 > 0$ ,*

$$\begin{aligned}
 \frac{dI_5}{dt} &\leq -\frac{kEh}{2} \|(w_{1x} - kw_3)\|^2 - \frac{\rho h}{2} \|w_{1t}\|^2 + k\rho h \|w_{3t}\|^2 \\
 &\quad + \frac{\rho h}{2} \|\phi_{2t}\|^2 + C(\varepsilon_5) \|\theta_{1xt}\|^2 + (kGh + \varepsilon_5) \|(\phi_2 + w_{3x} + kw_1)\|^2.
 \end{aligned}
 \tag{2.23}$$

*Proof.* Let (1) =  $-h\rho \int_0^1 w_{3t} (w_{1x} - kw_3) dx$ , (2) =  $-h\rho \int_0^1 w_{1t} (\phi_2 + w_{3x} + kw_1) dx$ , then using (1.1), (1.2), we have

$$\begin{aligned}
 (1)' &= -Gh \int_0^1 (\phi_2 + w_{3x} + kw_1)_x (w_{1x} - kw_3) dx - kEh \int_0^1 (w_{1x} - kw_3)^2 dx \\
 &\quad + \alpha k \int_0^1 \theta_{1t} (w_{1x} - kw_3) dx + k\rho h \int_0^1 w_{3t}^2 - \rho h \int_0^1 w_{3t} w_{1xt} dx, \\
 (2)' &= -Eh \int_0^1 (w_{1x} - kw_3)_x (\phi_2 + w_{3x} + kw_1) dx + \alpha \int_0^1 \theta_{1xt} (\phi_2 + w_{3x} + kw_1) dx \\
 &\quad + kGh \int_0^1 (\phi_2 + w_{3x} + kw_1)^2 dx - \rho h \int_0^1 w_{1t}^2 dx - \rho h \int_0^1 w_{1t} \phi_{2t} dx + \rho h \int_0^1 w_{1tx} w_{3t} dx.
 \end{aligned}
 \tag{2.24}$$

Then, noticing  $E = G$ , again, from the above two equalities and Young's inequality, we can obtain the assertion.

Next, we set

$$I_6 = -\rho h \int_0^1 w_{3t} w_3 dx - \rho h \int_0^1 w_{1t} w_1 dx. \quad (2.25)$$

□

**Lemma 2.8.** *Letting  $w_1, w_3, \phi_2, \theta_1, \theta_3$  be a solution of (1.1)–(1.5), then one has*

$$\frac{dI_6}{dt} \leq -\rho h (\|w_{3t}\|^2 + \|w_{1t}\|^2) + C\|\theta_{1xt}\|^2 + C\|\phi_{2x}\|^2. \quad (2.26)$$

*Proof.* Using (1.1), (1.2), we have

$$\begin{aligned} I_6' &= -\rho h \int_0^1 w_{3t}^2 dx - \rho h \int_0^1 w_{1t}^2 dx + Eh \int_0^1 (w_{1x} - kw_3)^2 dx \\ &\quad + Gh \int_0^1 (\phi_2 + w_{3x} + kw_1)(w_{3x} + kw_1) dx - \alpha \int_0^1 \theta_{1t}(w_{1x} - kw_3) dx. \end{aligned} \quad (2.27)$$

Noticing (2.3) and (2.4), we have that  $\exists C_1 > 0$  satisfy the following:

$$-\alpha \int_0^1 \theta_{1t}(w_{1x} - kw_3) dx \leq C_1 \|\theta_{1xt}\|^2 - Eh \|w_{1x} - kw_3\|^2. \quad (2.28)$$

Similarly,

$$\begin{aligned} &Gh \int_0^1 (\phi_2 + w_{3x} + kw_1)(w_{3x} + kw_1) dx \\ &= Gh \|\phi_2 + w_{3x} + kw_1\|^2 - Gh \int_0^1 (\phi_2 + w_{3x} + kw_1) \phi_2 dx \\ &\leq C_1 \|\phi_{2x}\|^2. \end{aligned} \quad (2.29)$$

Then, insert (2.28) and (2.29) into (2.27), and the assertion of the lemma follows.

Now, we set

$$I_7 = \rho c \int_0^1 \theta_{1t} \theta_1 dx + \frac{1}{2} \|\theta_{1x}\|^2. \quad (2.30)$$

□

**Lemma 2.9.** *Letting  $w_1, w_3, \phi_2, \theta_1, \theta_3$  be a solution of (1.1)–(1.5), then one has, for all  $\varepsilon_7 > 0$ ,*

$$\frac{dI_7}{dt} \leq -\frac{1}{2} \|\theta_{1x}\|^2 + \rho c \|\theta_{3xt}\|^2 + C(\varepsilon_7) (\|w_{1t}\|^2 + \|w_{3t}\|^2). \quad (2.31)$$

*Proof.* Using (1.5), we have

$$\frac{dI_7}{dt} = -\|\theta_{1x}\|^2 + \alpha T_0 \int_0^1 w_{1t}\theta_{1x} dx + \alpha T_0 k \int_0^1 w_{3t}\theta_{1x} dx + \rho c \|\theta_{1t}\|^2. \quad (2.32)$$

Then, using Young's and Poincaré's inequalities, we can obtain the assertion.

Now, letting  $N, N_1, N_2, N_3, N_4, N_5, N_6, N_7 > 0$ , we define the Lyapunov functional  $\mathcal{F}$  as follows:

$$\mathcal{F} = NE + N_1 I_1 + N_2 I_2 + N_3 I_3 + N_4 I_4 + N_5 I_5 + N_6 I_6 + N_7 I_7. \quad (2.33)$$

By using (2.4), (2.9), (2.13), (2.16), (2.19), (2.23), (2.26), and (2.31), we have

$$\begin{aligned} \frac{d\mathcal{F}}{dt} \leq & \Upsilon_1 \|\theta_{1xt}\|^2 + \Upsilon_2 \|\theta_{3x}\|^2 + \Upsilon_3 \|\phi_{2x}\|^2 + \Upsilon_4 \|\omega_{1t}\|^2 + \Upsilon_5 \|\phi_{2t}\|^2 \\ & + \Upsilon_6 \|\phi_2 + \omega_{3x} + k\omega_1\|^2 + \Upsilon_7 \|\omega_{1x} - k\omega_3\|^2 + \Upsilon_8 \|\omega_{3t}\|^2 + \Upsilon_9 \|\theta_{1x}\|^2, \end{aligned} \quad (2.34)$$

where

$$\begin{aligned} \Upsilon_1 &= -\frac{N}{T_0} + C(\varepsilon_1)N_1 + N_2 C(\varepsilon_2) + N_4 C(\varepsilon_4) + N_5 C(\varepsilon_5) + N_7 C_1 + \rho c N_7, \\ \Upsilon_2 &= -\frac{N}{T_0} + C(\varepsilon_1)N_1 + N_3 C(\varepsilon_3) + N_4 C(\varepsilon_4), \\ \Upsilon_3 &= -\frac{N_1 EI}{2} + \varepsilon_3 N_3 + C(\varepsilon_4)N_4 + C_1 N_6, \\ \Upsilon_4 &= -\frac{\alpha \rho h T_0 N_2}{2} + \frac{k h \rho I N_4}{2} - \frac{\rho h N_5}{2} - \rho h N_6 + N_7 C(\varepsilon_7), \\ \Upsilon_5 &= -\frac{\alpha \rho h T_0 N_3}{2} + \frac{k h \rho I N_4}{2} + N_1 \rho I + N_1 C(\varepsilon_1) + \frac{\rho h N_5}{2}, \\ \Upsilon_6 &= -\frac{G h^2 N_4}{2} + k G h N_5 + N_5 \varepsilon_5 + N_3 \varepsilon_3 + N_2 \varepsilon_2, \\ \Upsilon_7 &= -\frac{k E h N_5}{2} + N_4 \varepsilon_4 + N_1 \varepsilon_1 + N_2 \varepsilon_2, \\ \Upsilon_8 &= -N_6 \rho h + N_5 k \rho h + C(\varepsilon_2)N_2 + N_1 \varepsilon_1 + N_7 C(\varepsilon_7), \\ \Upsilon_9 &= -\frac{N_7}{2} + N_2 C(\varepsilon_2). \end{aligned} \quad (2.35)$$

We can choose  $N$  big enough,  $\varepsilon_1, \dots, \varepsilon_8$  small enough, and

$$\begin{aligned}
 N_1 &\gg N_4, N_6, \\
 N_2 &\gg N_4, \\
 N_3 &\gg N_1, N_6, \\
 N_4 &\gg N_5, \\
 N_6 &\gg N_2, N_5, N_7, \\
 N_7 &\gg N_2.
 \end{aligned} \tag{2.36}$$

Then  $\Upsilon_1, \dots, \Upsilon_9$  are all negative constants; at this point, there exists a constant  $\omega > 0$ , and (2.34) takes the form

$$\begin{aligned}
 \frac{d\mathcal{F}}{dt} \leq & -\omega \left( \|\theta_{1xt}\|^2 + \|\theta_{1x}\|^2 + \|\theta_{3x}\|^2 + \|\phi_{2x}\|^2 + \|w_{1t}\|^2 \right. \\
 & \left. + \|\phi_{2t}\|^2 + \|\phi_2 + w_{3x} + kw_1\|^2 + \|w_{1x} - kw_3\|^2 + \|w_{3t}\|^2 \right).
 \end{aligned} \tag{2.37}$$

We are now ready to prove Theorem 2.2. □

*Proof of Theorem 2.2.* Firstly, from the definition of  $\mathcal{F}$ , we have

$$\mathcal{F} \sim E(t), \tag{2.38}$$

which, from (2.37) and (2.38), leads to

$$\frac{d}{dt} \mathcal{F} \leq -\mu \mathcal{F}. \tag{2.39}$$

Integrating (2.39) over  $(0, t)$  and using (2.38) lead to (2.6). This completes the proof of Theorem 2.2. □

### 3. Global Attractors

In this section, we establish the existence of the global attractor for system (1.1)–(1.5).

Setting  $v = w_{1t}$ ,  $\varphi = w_{3t}$ ,  $\psi = \phi_{2t}$ ,  $\eta = \theta_{1t}$ , then, (1.1)–(1.5) can be transformed into the system

$$\begin{aligned}
 w_{1t} &= v, \\
 w_{3t} &= \varphi, \\
 \phi_{2t} &= \psi, \\
 \theta_{1t} &= \eta, \\
 \rho h v_t &= (Eh(w_{1x} - kw_3) - \alpha\theta_1)_x - kGh(\phi_2 + w_{3x} + kw_1), \\
 \rho h \varphi_t &= Gh(\phi_2 + w_{3x} + kw_1)_x + kEh(w_{1x} - kw_3) - k\alpha\theta_1, \\
 \rho I \psi_t &= EI\phi_{2xx} - Gh(\phi_2 + w_{3x} + kw_1) - \alpha\theta_{3x}, \\
 \rho c \eta_t &= \theta_{1xxt} + \theta_{1xx} - \alpha T_0(w_{1tx} - kw_{3t}), \\
 \rho c \theta_{3t} &= \theta_{3xx} - \alpha T_0 \phi_{2tx}.
 \end{aligned} \tag{3.1}$$

We consider the problem in the following Hilbert space:

$$\mathcal{H} = H_0^1 \times H_*^1 \times H_0^1 \times H_0^1 \times (L^2)^5. \tag{3.2}$$

Recall that the global attractor of  $S(t)$  acting on  $\mathcal{H}$  is a compact set  $\mathcal{A} \subset \mathcal{H}$  enjoying the following properties:

- (1)  $\mathcal{A}$  is fully invariant for  $S(t)$ , that is,  $S(t)\mathcal{A} = \mathcal{A}$  for every  $t \geq 0$ ;
- (2)  $\mathcal{A}$  is an attracting set, namely, for any bounded set  $\mathcal{R} \subset \mathcal{H}$ ,

$$\lim_{t \rightarrow \infty} \delta_{\mathcal{H}}(S(t)\mathcal{R}, \mathcal{A}) = 0, \tag{3.3}$$

where  $\delta_{\mathcal{H}}$  denotes the Hausdorff semidistance on  $\mathcal{H}$ .

More details on the subject can be found in the books [23, 26, 27].

*Remark 3.1.* The uniform energy estimate (2.6) implies the existence of a bounded absorbing set  $\mathcal{R}^* \subset \mathcal{H}$  for the  $C_0$  semigroup  $S(t)$ . Indeed, if  $\mathcal{R}^*$  is any ball of  $\mathcal{H}$ , then for any bounded set  $\mathcal{R} \subset \mathcal{H}$ , it is immediate to see that there exists  $t(\mathcal{R}) \geq 0$  such that

$$S(t)\mathcal{R} \subset \mathcal{R}^* \tag{3.4}$$

for every  $t \geq t(\mathcal{R})$ .

Moreover, if we define

$$\mathcal{R}_0 = \bigcup_{t \geq 0} S(t)\mathcal{R}^*, \quad (3.5)$$

it is clear that  $\mathcal{R}_0$  is still a bounded absorbing set which is also invariant for  $S(t)$ , that is,  $S(t)\mathcal{R}_0 \subset \mathcal{R}_0$  for every  $t \geq 0$ .

In the sequel, we define the operator  $A$  as  $Af = -f_{xx}$  with Dirichlet boundary conditions. It is well known that  $A$  is a positive operator on  $L^2$  with domain  $\mathfrak{D}(A) = H^2 \cap H_0^1$ . Moreover, we can define the powers  $A^s$  of  $A$  for  $s \in \mathbb{R}$ . The space  $V_{2s} = \mathfrak{D}(A^s)$  turns out to be a Hilbert space with the inner product

$$\langle u, v \rangle_{V_{2s}} = \langle A^s u, A^s v \rangle, \quad (3.6)$$

where  $\langle \cdot \rangle$  stands for  $L^2$ -inner product on  $L^2$ .

In particular,  $V_{-1} = H^{-1}$ ,  $V_0 = L^2$ , and  $V_1 = H_0^1$ . The injection  $V_{s_1} \hookrightarrow V_{s_2}$  is compact whenever  $s_1 > s_2$ . For further convenience, for  $s \in \mathbb{R}$ , introduce the Hilbert space

$$\mathcal{H}_s = V_{1+s} \times V_{1+s} \times V_{1+s} \times V_{1+s} \times (V_s)^5. \quad (3.7)$$

Clearly,  $\mathcal{H}_0 = \mathcal{H}$ .

Now, let  $z_0 = (u_0, w_0, \phi_0, \theta_0, v_0, \varphi_0, \psi_0, \eta_0, \xi_0)$ , where  $\mathcal{R}_0$  is the invariant, bounded absorbing set of  $S(t)$  given by Remark 3.1, and take the inner product in  $\mathcal{H}_0$  of (3.1) and  $(A^\sigma w_1, A^\sigma w_3, A^\sigma \phi_2, A^\sigma \theta_1, A^\sigma v, A^\sigma \varphi, A^\sigma \psi, A^\sigma \eta, A^\sigma \theta_3)$  to get

$$\begin{aligned} & \frac{d}{dt} \left( Eh \| (w_{1x} - kw_3) \|_\sigma^2 + Gh \| (\phi_2 + w_{3x} + kw_1) \|_\sigma^2 + EI \| \phi_2 \|_{1+\sigma}^2 \right. \\ & \quad \left. + \rho h \left( \| w_{1t} \|_\sigma^2 + \| w_{3t} \|_\sigma^2 \right) + \rho I \| \phi_{2t} \|_\sigma^2 + \frac{\rho C}{T_0} \left( \| \theta_{1t} \|_\sigma^2 + \| \theta_1 \|_{1+\sigma}^2 + \| \theta_3 \|_\sigma^2 \right) \right) \\ & = -\frac{2}{T_0} \left( \| \theta_{1t} \|_{1+\sigma}^2 + \| \theta_3 \|_{(1+\sigma)}^2 \right). \end{aligned} \quad (3.8)$$

Here, the boundary term of integration by parts is neglected since we are working with more regular functions. We denote

$$\begin{aligned}
 E_2(t) &= Eh\|(w_{1x} - kw_3)\|_\sigma^2 + Gh\|(\phi_2 + w_{3x} + kw_1)\|_\sigma^2 + EI\|\phi_2\|_{1+\sigma}^2 \\
 &\quad + \rho h(\|w_{1t}\|_\sigma^2 + \|w_{3t}\|_\sigma^2) + \rho I\|\phi_{2t}\|_\sigma^2 + \frac{\rho c}{T_0}(\|\theta_1\|_\sigma^2 + \|\theta_3\|_\sigma^2), \\
 F_1(t) &= \int_0^1 \rho I A^\sigma \phi_{2t} \phi_2 dx + \int_0^1 \rho h A^\sigma w_{3t} f dx, \\
 F_2(t) &= \rho c \rho h \int_0^1 \left( \int_0^x \theta_1 dy \right) A^\sigma w_{1t} dx, \\
 F_3 &= \rho c \rho I \int_0^1 \left( \int_0^x \theta_3 dy \right) A^\sigma \phi_{2t} dx, \\
 F_4(t) &= h \rho I \int_0^1 A^\sigma \phi_{2t} (\phi_2 + w_{3x} + kw_1) dx + h \rho I \int_0^1 A^\sigma \phi_{2x} w_{3t} dx, \\
 F_5(t) &= -h \rho \int_0^1 A^\sigma w_{3t} (w_{1x} - kw_3) dx - h \rho \int_0^1 A^\sigma w_{1t} (\phi_2 + w_{3x} + kw_1) dx, \\
 F_6(t) &= -\rho h \int_0^1 A^\sigma w_{3t} w_3 dx - \rho h \int_0^1 A^\sigma w_{1t} w_1 dx, \\
 F_7 &= \rho c \int_0^1 A^\sigma \theta_{1t} \theta_1 dx + \frac{1}{2} \int_0^1 A^\sigma \theta_{1x}^2 dx.
 \end{aligned} \tag{3.9}$$

Then, introduce the functional

$$\mathcal{J}(t) = NE_2(t) + N_1F_1 + N_2F_2 + N_3F_3 + N_4F_4 + N_5F_5 + N_6F_6 + N_7F_7. \tag{3.10}$$

By repeating similar argument as in the proofs of Lemmas 2.3–2.9 and (3.8), choosing our constants very carefully and properly, we get

$$\frac{d}{dt} \mathcal{J}(t) + cE_2(t) \leq 0. \tag{3.11}$$

On the other hand,

$$\mathcal{J}(t) \sim E_2(t), \tag{3.12}$$

so that

$$\frac{d}{dt} \mathcal{J}(t) + c_1 \mathcal{J}(t) \leq 0, \tag{3.13}$$

which gives

$$E_2(t) \sim \mathcal{J}(t) \leq c_2 e^{-c_1 t}, \quad \|z(t)\|_{\mathcal{L}_\sigma} \leq c_2 e^{-c_1 t}. \quad (3.14)$$

Let  $\mathcal{R}(t)$  be the ball of  $V_{3/2} \times V_{3/2} \times V_{3/2} \times V_{3/2} \times (V_{1/2})^5$ ; from the compact embedding  $V_{3/2} \times V_{3/2} \times V_{3/2} \times V_{3/2} \times (V_{1/2})^5 \hookrightarrow H_0^1 \times H_*^1 \times H_0^1 \times H_0^1 \times (L^2)^5$ ,  $\mathcal{R}(t)$  is compact in  $\mathcal{L}$ . Then, due to the compactness of  $\mathcal{R}(t)$ , for every fixed  $t \geq 0$  and every  $d > c_2 e^{-c_1 t}$ , there exist finitely many balls of  $\mathcal{L}$  of radius  $d$  such that  $z(t)$  belongs to the union of such balls, for every  $z_0 \in \mathcal{R}_0$ . This implies that

$$\alpha_{\mathcal{L}}(S(t)\mathcal{R}_0) \leq c_2 e^{-c_1 t}, \quad \forall t \geq 0, \quad (3.15)$$

where  $\alpha_{\mathcal{L}}$  is the Kuratowski measure of noncompactness, defined by

$$\alpha_{\mathcal{L}}(\mathcal{R}) = \inf\{d : \mathcal{R} \text{ has a finite cover of balls of } \mathcal{L} \text{ of diameter less than } d\}. \quad (3.16)$$

Since the invariant, connected, bounded absorbing set  $\mathcal{R}_0$  fulfills (3.15), exploiting a classical result of the theory of attractors of semigroups (see, e.g., [28]), we conclude that the  $\omega$ -limit set of  $\mathcal{R}_0$ , that is,

$$\mathcal{A} \equiv \omega(\mathcal{R}_0) = \bigcap_{t \geq 0} \overline{\bigcup_{s \geq t} S(s)\mathcal{R}_0}, \quad (3.17)$$

is a connected and compact global attractor of  $S(t)$ . Therefore, we have proved the following result.

**Theorem 3.2.** *Under the assumption of  $(H_1) - (H_2)$ , problem (3.1) possesses a unique global attractor  $\mathcal{A}$ .*

## References

- [1] J. A. C. Bresse, *Cours de Mécanique Appliquée*, Mallet Bachelier, Paris, France, 1859.
- [2] S. A. Messaoudi and B. Said-Houari, "Energy decay in a Timoshenko-type system of thermoelasticity of type III," *Journal of Mathematical Analysis and Applications*, vol. 348, no. 1, pp. 298–307, 2008.
- [3] S. A. Messaoudi and B. Said-Houari, "Energy decay in a Timoshenko-type system with history in thermoelasticity of type III," *Advances in Differential Equations*, vol. 14, no. 3-4, pp. 375–400, 2009.
- [4] F. Ammar-Khodja, A. Benabdallah, J. E. Muñoz Rivera, and R. Racke, "Energy decay for Timoshenko systems of memory type," *Journal of Differential Equations*, vol. 194, no. 1, pp. 82–115, 2003.
- [5] J. E. Muñoz Rivera and H. D. Fernández Sare, "Stability of Timoshenko systems with past history," *Journal of Mathematical Analysis and Applications*, vol. 339, no. 1, pp. 482–502, 2008.
- [6] H. D. Fernández Sare and R. Racke, "On the stability of damped Timoshenko systems: Cattaneo versus Fourier law," *Archive for Rational Mechanics and Analysis*, vol. 194, no. 1, pp. 221–251, 2009.
- [7] K. Liu, Z. Liu, and B. Rao, "Exponential stability of an abstract nondissipative linear system," *SIAM Journal on Control and Optimization*, vol. 40, no. 1, pp. 149–165, 2001.
- [8] S. A. Messaoudi, M. Pokojovy, and B. Said-Houari, "Nonlinear damped Timoshenko systems with second sound—global existence and exponential stability," *Mathematical Methods in the Applied Sciences*, vol. 32, no. 5, pp. 505–534, 2009.

- [9] J. E. Muñoz Rivera and R. Racke, "Mildly dissipative nonlinear Timoshenko systems—global existence and exponential stability," *Journal of Mathematical Analysis and Applications*, vol. 276, no. 1, pp. 248–278, 2002.
- [10] A. Soufyane, "Stabilisation de la poutre de Timoshenko," *Comptes Rendus de l'Académie des Sciences. Série I*, vol. 328, no. 8, pp. 731–734, 1999.
- [11] G. Avalos and I. Lasiecka, "Exponential stability of a thermoelastic system with free boundary conditions without mechanical dissipation," *SIAM Journal on Mathematical Analysis*, vol. 29, no. 1, pp. 155–182, 1998.
- [12] G. Avalos and I. Lasiecka, "Exponential stability of a thermoelastic system without mechanical dissipation," *Rendiconti dell'Istituto di Matematica dell'Università di Trieste*, vol. 28, pp. 1–28, 1996.
- [13] I. Chueshov and I. Lasiecka, "Inertial manifolds for von Kármán plate equations," *Applied Mathematics and Optimization*, vol. 46, no. 2-3, pp. 179–206, 2002.
- [14] I. Chueshov and I. Lasiecka, "Long-time behavior of second order evolution equations with nonlinear damping," *Memoirs of the American Mathematical Society*, vol. 195, no. 912, 2008.
- [15] I. Chueshov and I. Lasiecka, *Von Karman Evolution Equations: Well-posedness and Long Time Dynamics*, Springer Monographs in Mathematics, Springer, New York, NY, USA, 2010.
- [16] Z. Liu and B. Rao, "Energy decay rate of the thermoelastic Bresse system," *Zeitschrift für Angewandte Mathematik und Physik*, vol. 60, no. 1, pp. 54–69, 2009.
- [17] J. Prüss, "On the spectrum of  $C_0$ -semigroups," *Transactions of the American Mathematical Society*, vol. 284, no. 2, pp. 847–857, 1984.
- [18] F. L. Huang, "Characteristic conditions for exponential stability of linear dynamical systems in Hilbert spaces," *Annals of Differential Equations*, vol. 1, no. 1, pp. 43–56, 1985.
- [19] S. Borini and V. Pata, "Uniform attractors for a strongly damped wave equation with linear memory," *Asymptotic Analysis*, vol. 20, no. 3-4, pp. 263–277, 1999.
- [20] I. Chueshov and I. Lasiecka, "Attractors and long time behavior of von Karman thermoelastic plates," *Applied Mathematics and Optimization*, vol. 58, no. 2, pp. 195–241, 2008.
- [21] C. Giorgi, J. E. Muñoz Rivera, and V. Pata, "Global attractors for a semilinear hyperbolic equation in viscoelasticity," *Journal of Mathematical Analysis and Applications*, vol. 260, no. 1, pp. 83–99, 2001.
- [22] V. Pata and A. Zucchi, "Attractors for a damped hyperbolic equation with linear memory," *Advances in Mathematical Sciences and Applications*, vol. 11, no. 2, pp. 505–529, 2001.
- [23] Y. Qin, *Nonlinear Parabolic-Hyperbolic Coupled Systems and Their Attractors*, vol. 184 of *Operator Theory: Advances and Applications*, Birkhäuser, Basel, Switzerland, 2008.
- [24] S. Zheng and Y. Qin, "Maximal attractor for the system of one-dimensional polytropic viscous ideal gas," *Quarterly of Applied Mathematics*, vol. 59, no. 3, pp. 579–599, 2001.
- [25] Z. Liu and S. Zheng, *Semigroups Associated with Dissipative Systems*, vol. 398 of *Chapman & Hall/CRC Research Notes in Mathematics*, Chapman & Hall/CRC, Boca Raton, Fla, USA, 1999.
- [26] A. Pazy, *Semigroups of Linear Operators and Applications to Partial Differential Equations*, vol. 44 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1983.
- [27] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, vol. 68 of *Applied Mathematical Sciences*, Springer, New York, NY, USA, 1988.
- [28] J. K. Hale, "Asymptotic behaviour and dynamics in infinite dimensions," in *Nonlinear Differential Equations*, J. Hale and P. Martinez-Amores, Eds., vol. 132, pp. 1–42, Pitman, Boston, Mass, USA, 1985.