

RESEARCH

Open Access

# Local bifurcation of limit cycles and integrability of a class of nilpotent systems

Hongwei Li<sup>1</sup>, Yusen Wu<sup>2</sup> and Yinlai Jin<sup>1\*</sup>

\* Correspondence: [jinyinlai@lyu.edu.cn](mailto:jinyinlai@lyu.edu.cn)

<sup>1</sup>School of Science, Linyi University, Linyi 276005, Shandong, P.R. China  
Full list of author information is available at the end of the article

## Abstract

In this article, center conditions and bifurcation of limit cycles at the nilpotent critical point in a class of seventh-degree systems are investigated. With the help of computer algebra system MATHEMATICA, the first 13 quasi-Lyapunov constants are deduced. As a result, sufficient and necessary conditions in order to have a center are obtained. The result that there exist 13 small amplitude limit cycles created from the three-order nilpotent critical point is also proved. Henceforth, we give a lower bound of cyclicity of three-order nilpotent critical point for seventh-degree nilpotent systems.

**MSC:** 34C05; 34C07.

**Keywords:** three-order nilpotent critical point, center-focus problem, bifurcation of limit cycles, quasi-Lyapunov constant

## 1 Introduction

Computation of Lyapunov quantities is a hot topic with large number of articles per year, but it is very difficult to obtain general results. The methods of computation of Lyapunov quantities when the critical points are non-degenerate have been greatly developed by many mathematicians. The method in [1,2] is based on the sequential construction of Lyapunov functions. Furthermore, computing Lyapunov quantities using the reduction of system to normal form could be seen in [3-5]. Another approach to numerical computation of Lyapunov quantities which uses the passage to the polar coordinates and the procedure of sequential construction of solution approximations is related with the obtaining of approximations of system solution, see [2], they also could be seen in [6,7]. But computations of Lyapunov quantities become difficult when the critical points are degenerate because the method of the Poincaré formal series cannot be used in order to compute Lyapunov constants in a neighborhood of the critical point.

The nilpotent center problem was investigated by Moussu [8] and Stróżyńska and Żołądek [9]. In [10], Takens proved that Lyapunov system can be formally transformed into a generalized Liénard system. Furthermore, in [11], Álvarez and Gasull proved that the generalized Liénard system could be simplified even more by a reparametrization of the time. At the same time, Giacomini et al. [12,13] proved that the analytic nilpotent systems with a center can be expressed as limit of non-degenerate systems with a center.

As far as we know, there are essentially three differential ways of obtaining Lyapunov constant for nilpotent critical points in theory: by using normal form theory [14], by computing the Poincaré return map [15] or by using Lyapunov functions [16]. Álvarez investigated the momodromy and stability for nilpotent critical points with the method of computing the Poincaré return map, see for instance [17]; Chavarriga et al. investigated the local analytic integrability for nilpotent centers by using Lyapunov functions, see for instance [18]; Moussu investigated the center-focus problem of nilpotent critical points with the method of normal form theory, see for instance [8].

Nevertheless, these methods are so complicated that it is hard to use for an analytic system with a monodromic point, even in the case of a concrete polynomial systems. So there are very few results known for concrete differential systems with monodromic nilpotent critical points. Gasull and Torregrosa [19] have generalized the scheme of computation of Lyapunov constants for systems of the form

$$\begin{aligned} \dot{x} &= y + \sum_{k \geq n+1} F_k(x, y), \\ \dot{y} &= -x^{2n-1} + \sum_{k \geq 2n} G_k(x, y), \end{aligned} \tag{1.1}$$

where  $F_k$  and  $G_k$  are  $(1, n)$ -quasi-homogeneous functions of degree  $k$ . Chavarriga et al. investigated the integrability of centers perturbed by  $(p, q)$ -quasi-homogeneous polynomials in [20]. Fortunately, Yirong Liu and Jibin Li [21] found that there always exists a formal inverse integrating factor for three-order nilpotent critical points in 2009, and they gave a new definition of the focal values under the generalized triangle polar coordinates and the method of commuting Lyapunov constants using the inverse integral factors for the three-order nilpotent critical point.

For a given family of polynomial differential equations, the number of Lyapunov constants needed to solve the center-focus problem is also related with the so-called cyclicity of the point, i.e., the number of limit cycles that appear from it by small perturbations of the coefficients of the given differential equation inside the family considered (see, [22] for cases where this relation does not exist for the case of non-degenerate centers). Let  $N(n)$  be the maximum possible number of limit cycles bifurcating from nilpotent critical points for analytic vector fields of degree  $n$ . It was found that  $N(3) \geq 2$ ,  $N(5) \geq 5$ ,  $N(7) \geq 9$  in [23],  $N(3) \geq 3$ ,  $N(5) \geq 5$  in [17], and for Kukles system with six parameters  $N(3) \geq 3$  in [11]. Recently, Yirong Liu and Jibin Li proved that  $N(3) \geq 8$  in [24]. In this article, by employing the inverse integral factor method introduced in [21], we consider a planar septic ordinary differential equation having a three-order nilpotent critical point with the form

$$\begin{aligned} \frac{dx}{dt} &= \mu y + \mu x^3 - \mu x^2 y + a_{12} x y^2 + a_{03} y^3 + \left(1 - \frac{71275\mu}{378}\right) x^4 y + a_{32} x^3 y^2 \\ &\quad + a_{23} x^2 y^3 + a_{14} x y^4 + a_{05} y^5 - \mu y(x^2 + y^2)^3, \\ \frac{dy}{dt} &= -2\mu x^3 + b_{21} x^2 y + \mu x y^2 + b_{03} y^3 - 2 \left(1 - \frac{71275\mu}{378}\right) x^3 y^2 + b_{23} x^2 y^3 \\ &\quad + b_{14} x y^4 + b_{05} y^5 + \mu x(x^2 + y^2)^3. \end{aligned} \tag{1.2}$$

We will prove  $N(7) \geq 13$ . To the best of authors' knowledge, their result on the lower bounds of cyclicity of three-order nilpotent critical points for septic systems is new. It is helpful to Hilbert's 16th problem.

The rest of the article is organized as follows. In Section 2, some preliminary knowledge given in [21] which is useful throughout the article are introduced. In Section 3, using the linear recursive formulae in [21] to do direct computation, the first 13 quasi-Lyapunov constants and the sufficient and necessary conditions of center are obtained. This article is ended with Section 4 in which the 13-order weak focus conditions and the result that there exist 13 limit cycles in the neighborhood of the three-order nilpotent critical point is proved.

## 2 Preliminary knowledge

The idea of this section comes from [21,24], where the center-focus problem of three-order nilpotent critical points of the planar dynamical systems is studied. For more details, please refer to [21,24]. We will recall the related notions and results. The origin of system

$$\begin{aligned} \frac{dx}{dt} &= \gamma + \sum_{i+j=2}^{\infty} a_{ij}x^i\gamma^j = X(x, \gamma), \\ \frac{dy}{dt} &= \sum_{i+j=2}^{\infty} b_{ij}x^i\gamma^j = Y(x, \gamma). \end{aligned} \tag{2.1}$$

is a three-order monodromic critical point if and only if the system could be written as follows:

$$\begin{aligned} \frac{dx}{dt} &= \gamma + \mu x^2 + \sum_{i+2j=3}^{\infty} a_{ij}x^i\gamma^j = X(x, \gamma), \\ \frac{dy}{dt} &= -2x^3 + 2\mu x\gamma + \sum_{i+2j=4}^{\infty} b_{ij}x^i\gamma^j = Y(x, \gamma). \end{aligned} \tag{2.2}$$

Under the transformation of generalized polar coordinates

$$x = r \cos \theta, \gamma = r^2 \sin \theta, \tag{2.3}$$

system (2.2) can be changed into

$$\frac{dr}{d\theta} = \frac{-\cos \theta [\sin \theta (1 - 2\cos^2 \theta) + \mu (\cos^2 \theta + 2\sin^2 \theta)]}{2(\cos^4 \theta + \sin^2 \theta)} r + o(r). \tag{2.4}$$

In a small neighborhood, we can define the successor function of system (2.2) as follows:

$$\Delta(h) = \tilde{r}(-2\pi, h) - h = \sum_{k=2}^{\infty} v_k(-2\pi)h^k. \tag{2.5}$$

We have the following result.

**Lemma 2.1.** For any positive integer  $m$ ,  $v_{2m+1}(-2\pi)$  has the form

$$v_{2m+1}(-2\pi) = \sum_{k=1}^m S_k^{(m)} v_{2k}(-2\pi), \tag{2.6}$$

where  $S_k^{(m)}$  is a polynomial of  $v_j(\pi)$ ,  $v_j(2\pi)$ ,  $v_j(-2\pi)$ , ( $j = 2, 3, \dots, 2m$ ) with rational coefficients.

It is different from the center-focus problem for the elementary critical points, we know from Lemma 2.1 that when  $k > 1$  for the first non-zero  $v_k(-2\pi)$ ,  $k$  is an even integer.

**Definition 2.1.** 1. For any positive integer  $m$ ,  $v_{2m}(-2\pi)$  is called the  $m$ -order focal value of system (2.2) in the origin.

2. If  $v_2(-2\pi) \neq 0$ , the origin of system (2.2) is called 1-order weak focus. If there is an integer  $m > 1$ , such that  $v_2(-2\pi) = v_4(-2\pi) = \dots = v_{2m-2}(-2\pi) = 0$ ,  $v_{2m}(-2\pi) \neq 0$ , then, the origin of system (2.2) is called  $m$ -order weak focus.

3. If for all positive integer  $m$ , we have  $v_{2m}(-2\pi) = 0$ , then, the origin of system (2.2) is called a center.

Consider the system

$$\begin{aligned} \frac{dx}{dt} &= \delta x + \gamma + \sum_{k+j=2}^{\infty} a_{kj}(\gamma) x^k \gamma^j, \\ \frac{dy}{dt} &= 2\delta y + \sum_{k+j=2}^{\infty} b_{kj}(\gamma) x^k \gamma^j, \end{aligned} \tag{2.7}$$

where  $\gamma = \{\gamma_1, \gamma_2, \dots, \gamma_{m-1}\}$  is  $(m-1)$ -dimensional parameter vector. Let  $\gamma_0 = \{\gamma_1^{(0)}, \gamma_2^{(0)}, \dots, \gamma_{m-1}^{(0)}\}$  be a point at the parameter space. Suppose that for  $\|\gamma - \gamma_0\| \ll 1$ , the functions of the right hand of system (2.7) are power series of  $x, y$  with a non-zero convergence radius and have continuous partial derivatives with respect to  $\gamma$ . In addition,

$$a_{20}(\gamma) \equiv \mu, b_{20}(\gamma) \equiv 0, b_{11}(\gamma) \equiv 2\mu, b_{30}(\gamma) \equiv -2. \tag{2.8}$$

For an integer  $k$ , letting  $v_{2k}(-2\pi, \gamma)$  be the  $k$ -order focal value of the origin of system (2.7) <sub>$\delta = 0$</sub> .

**Theorem 2.1.** If for  $\gamma = \gamma_0$ , the origin of system (2.7) <sub>$\delta = 0$</sub>  is a  $m$ -order weak focus, and the Jacobian

$$\frac{\partial(v_2, v_4, \dots, v_{2m-2})}{\partial(\gamma_1, \gamma_2, \dots, \gamma_{m-1})} \Big|_{\gamma=\gamma_0} \neq 0, \tag{2.9}$$

then, there exist two positive numbers  $\delta^*$  and  $\gamma^*$ , such that for  $0 < |\delta| < \delta^*$ ,  $0 < \|\gamma - \gamma_0\| < \gamma^*$ , in a neighborhood of the origin, system (2.7) has at most  $m$  limit cycles which enclose the origin (an elementary node)  $O(0,0)$ . In addition, under the above conditions, there exist  $\tilde{\gamma}, \tilde{\delta}$ , such that when  $\gamma = \tilde{\gamma}, \delta = \tilde{\delta}$ , there exist exact  $m$  limit cycles of (2.7) in a small neighborhood of the origin.

The following key results which define the quasi-Lyapunov constants and provide a way of computing them were also given by Liu and Li [24].

**Theorem 2.2.** *If the origin of system (2.2) is a s-class or ∞-class, one can construct successively the terms of the formal power series  $M(x, y) = x^4 + y^2 + o(r^4)$ , such that*

$$\frac{\partial}{\partial x} \left( \frac{X}{M^{s+1}} \right) + \frac{\partial}{\partial y} \left( \frac{Y}{M^{s+1}} \right) = \frac{1}{M^{s+2}} \sum_{m=1}^{\infty} (2m - 4s - 1) \lambda_m [x^{2m+4} + o(r^{2m+4})], \quad (2.10)$$

**Theorem 2.3.** *For system (2.2), if there exists a natural number s and a formal series  $M(x, y) = x^4 + y^2 + o(r^4)$ , such that (2.10) holds, then*

$$\{v_{2m}(-2\pi)\} \sim [\sigma_m(s, \mu) \lambda_m], \quad (2.11)$$

where

$$\sigma_m(s, \mu) = \frac{1}{2} \int_0^{2\pi} \frac{(1 + \sin^2 \theta) \cos^{2m+4} \theta}{(\cos^4 \theta + \sin^2 \theta)^{s+2}} v_1^{2m-4s-1}(\theta) d\theta. \quad (2.12)$$

**Definition 2.2.** *If there exists a natural number s and a formal series  $M(x, y) = x^4 + y^2 + o(r^4)$ , such that (2.10) holds, then,  $\lambda_m$  is called the m-th quasi-Lyapunov constants of the origin of system (2.2).*

**Theorem 2.4.** *For any positive integer s and a given number sequence*

$$\{c_{0\beta}\}, \beta \geq 3, \quad (2.13)$$

*one can construct successively the terms with the coefficients  $c_{\alpha\beta}$  satisfying  $\alpha \neq 0$  of the formal series*

$$M(x, y) = y^2 + \sum_{\alpha+\beta=3}^{\infty} c_{\alpha\beta} x^\alpha y^\beta = \sum_{k=2}^{\infty} M_k(x, y), \quad (2.14)$$

such that

$$\left( \frac{\partial X}{\partial x} + \frac{\partial Y}{\partial y} \right) M - (s + 1) \left( \frac{\partial M}{\partial x} X + \frac{\partial M}{\partial y} Y \right) = \sum_{m=3}^{\infty} \omega_m(s, \mu) x^m. \quad (2.15)$$

where for all k,  $M_k(x, y)$  is a k-homogeneous polynomial of x, y and  $\mu = 0$ .

It is easy to see that (2.15) is linear with respect to the function M, so that we can easily find the following recursive formulae for the calculation of  $c_{\alpha\beta}$  and  $\omega_m(s, \mu)$ .

**Theorem 2.5.** *For  $\alpha \geq 1, \alpha + \beta \geq 3$  in (2.14) and (2.15),  $c_{\alpha\beta}$  can be uniquely determined by the recursive formula*

$$c_{\alpha\beta} = \frac{1}{(s + 1)\alpha} (A_{\alpha-1, \beta+1} + B_{\alpha-1, \beta+1}). \quad (2.16)$$

For  $m \geq 1, \omega_m(s, \mu)$  can be uniquely determined by the recursive formula

$$\omega_m(s, \mu) = A_{m,0} + B_{m,0}, \quad (2.17)$$

where

$$A_{\alpha\beta} = \sum_{k+j=2}^{\alpha+\beta-1} [k - (s + 1)(\alpha - k + 1)] a_{kj} c_{\alpha-k, \beta-j},$$

$$B_{\alpha\beta} = \sum_{k+j=2}^{\alpha+\beta-1} [j - (s + 1)(\beta - j + 1)] b_{kj} c_{\alpha-k, \beta-j+1}. \quad (2.18)$$

Notice that in (2.18), we set

$$\begin{aligned} c_{00} &= c_{10} = c_{01} = 0, \\ c_{20} &= c_{11} = 0, c_{02} = 1, \\ c_{\alpha\beta} &= 0, \text{ if } \alpha < 0 \text{ or } \beta < 0. \end{aligned} \tag{2.19}$$

We see from Theorem 2.2 that if the origin of system (2.2) is  $s$ -class or  $\infty$ -class, then, by choosing  $\{c_{\alpha\beta}\}$ , such that

$$\omega_{2k+1}(s, \mu) = 0, k = 1, 2, \dots, \tag{2.20}$$

we can obtain a solution group of  $\{c_{\alpha\beta}\}$  of (2.20), thus, we have

$$\lambda_m = \frac{\omega_{2m+4}(s, \mu)}{2m - 4s - 1}. \tag{2.21}$$

Clearly, the recursive formulae by Theorem 2.5 is linear with respect to all  $c_{\alpha\beta}$ . Therefore, it is convenient to perform the computations by using computer algebraic system like MATHEMATICA.

### 3 Quasi-Lyapunov constants and center conditions

Theorem 2.4 implies that we can find a positive integer  $s$  and a formal series  $M(x, y) = x^4 + y^2 + o(r^4)$  for system (1.2) such that (2.15) holds. Meanwhile, with the help of MATH-EMATICA, by applying the recursive formulae presented in Theorem 2.5 to carry out calculations, we have

$$\begin{aligned} \omega_3 &= \omega_4 = \omega_5 = 0, \\ \omega_6 &= -\frac{1}{3\mu}(b_{21} + 3\mu)(-1 + 4s), \\ \omega_7 &= 3(s + 1)c_{03}, \\ \omega_8 &= -\frac{2}{5\mu}(a_{12} + 3b_{03})(-3 + 4s). \end{aligned} \tag{3.1}$$

(2.21) and (3.1) implies that the first two quasi-Lyapunov constants of system (1.2):

$$\begin{aligned} \lambda_1 &= \frac{\omega_6}{1 - 4s} = -\frac{1}{3\mu}(b_{21} + 3\mu), \\ \lambda_2 &= \frac{\omega_8}{3 - 4s} = \frac{2}{5\mu}(a_{12} + 3b_{03}). \end{aligned} \tag{3.2}$$

we see from  $\omega_7 = 0$  that

$$c_{03} = 0. \tag{3.3}$$

Furthermore, the following conclusion holds.

**Proposition 3.1.** For system (1.2), one can determine successively the terms of the formal series  $M(x, y) = x^4 + y^2 + o(r^4)$ , such that

$$\left(\frac{\partial X}{\partial x} + \frac{\partial Y}{\partial y}\right)M - 2\left(\frac{\partial M}{\partial x}X + \frac{\partial M}{\partial y}Y\right) = \sum_{m=1}^{13} \lambda_m [(2m - 5)x^{2m+4} + o(r^{30})], \tag{3.4}$$

where  $\lambda_m$  is the  $m$ th quasi-Lyapunov constant at the origin of system (1.2),  $m = 1, 2, \dots, 13$ .

**Theorem 3.1.** For system (1.2), the first 13 quasi-Lyapunov constants at the origin are given by

$$\begin{aligned}
 \lambda_1 &= -\frac{1}{3\mu}(b_{21} + 3\mu), \\
 \lambda_2 &= \frac{2}{5\mu}(a_{12} + 3b_{03}), \\
 \lambda_3 &= \frac{2}{7\mu}(a_{32} + b_{23}), \\
 \lambda_4 &= \frac{4}{15\mu}(a_{14} - a_{23} + 5b_{05} - 2b_{14}), \\
 \lambda_5 &= -\frac{40}{77\mu^2}(a_{23} + 2b_{14})(-b_{03} + \mu), \\
 \lambda_6 &= -\frac{2}{7371\mu^2}(a_{23} + 2b_{14})(-756 + 252a_{32} + 142739\mu), \\
 \lambda_7 &= -\frac{8}{6237\mu^2}(a_{23} + 2b_{14})(189 - 378b_{05} + 189b_{14} - 35516\mu), \\
 \lambda_8 &= -\frac{8}{125307\mu^2}(a_{23} + 2b_{14})(-945 + 2646a_{03} + 231391\mu), \\
 \lambda_9 &= -\frac{1}{358435\mu^2}(a_{23} + 2b_{14})(-34020 + 4116a_{23} + 1372b_{14} + 5024095\mu), \\
 \lambda_{10} &= \frac{1}{386822709\mu^3}(a_{23} + 2b_{14}) - 714420 - 594712314\mu - 14002632a_{05}\mu + 90016920a_{23}\mu \\
 &\quad + 102674821303\mu^2), \\
 \lambda_{11} &= -\frac{17}{6544091169\mu^3}(a_{23} + 2b_{14})(-359281818 + 42007896a_{23} + 36356476968\mu + 2196158788177\mu^2), \\
 \lambda_{12} &= -\frac{1}{35823651080490\mu^4}(a_{23} + 2b_{14})(-26843045544 + 15035266598814\mu - 2015039855231046\mu^2 \\
 &\quad - 154659205390636303\mu^3 + 30287803437510297551\mu^4), \\
 \lambda_{13} &= -\frac{1}{1549838616695238870\mu^4}(a_{23} + 2b_{14})(-153852012319592664 + 34933988751683499324\mu \\
 &\quad + 10292695776995494843134\mu^2 - 3251791005302797375353453\mu^3 \\
 &\quad + 218543335369252401019315591\mu^4).
 \end{aligned} \tag{3.5}$$

In the above expressions of  $\lambda_k$ , we have already, let  $\lambda_1 = \lambda_2 = \dots = \lambda_{k-1} = 0$ ,  $k = 2, \dots, 13$ .

It follows from Theorem 3.1 that

**Proposition 3.2.** The first 13 quasi-Lyapunov constants at the origin of system (1.2) are zero if and only if the following condition is satisfied:

$$b_{21} = -3\mu, \quad a_{12} = -3b_{03}, \quad b_{23} = -a_{32}, \quad a_{14} = -5b_{05}, \quad a_{23} = -2b_{14}.$$

When the condition of Proposition 3.2 holds, system (1.2) can be brought to

$$\begin{aligned}
 \frac{dx}{dt} &= \mu y + \mu x^3 - \mu x^2 y - 3b_{03}xy^2 + a_{03}y^3 + \left(1 - \frac{71275\mu}{378}\right)x^4 y + a_{32}x^3 y^2 \\
 &\quad - 2b_{14}x^2 y^3 - 5b_{05}xy^4 + a_{05}y^5 - \mu y(x^2 + y^2)^3, \\
 \frac{dy}{dt} &= -2\mu x^3 - 3\mu x^2 y + \mu xy^2 + b_{03}y^3 - 2\left(1 - \frac{71275\mu}{378}\right)x^3 y^2 - a_{32}x^2 y^3 \\
 &\quad + b_{14}xy^4 + b_{05}y^5 + \mu x(x^2 + y^2)^3.
 \end{aligned} \tag{3.6}$$

system (3.6) has an analytic first integral

$$\begin{aligned}
 H(x, y) &= \frac{\mu}{2}y^2 + \frac{\mu}{2}x^4 + \frac{a_{03}}{4}y^4 + \frac{a_{05}}{5}y^5 - b_{05}xy^5 - b_{03}xy^3 - \frac{\mu}{2}x^2 y^2 + \mu x^3 y \\
 &\quad + \frac{1}{2}\left(1 - \frac{71275\mu}{378}\right)x^4 y^2 - \frac{b_{14}}{2}x^2 y^4 + \frac{a_{32}}{3}x^3 y^3 - \frac{\mu}{8}(x^2 + y^2)^4
 \end{aligned} \tag{3.7}$$

Proposition 3.2 implies that

**Proposition 3.3.** The origin of systems (3.6) is a center.

From Propositions 3.2 and 3.3, we further have

**Theorem 3.2.** *The origin of system (1.2) is a center if and only if the first 13 quasi-Lyapunov constants are zero, that is, the condition in Proposition 3.2 is satisfied.*

**4 Multiple bifurcation of limit cycles**

It is very interesting to study the number of limit cycles which could be bifurcated from critical point  $O(0,0)$ , because it is closely related with 16th problem of 23 problems of Hilbert. In this section, we will prove that the perturbed system of (1.2) can generate 13 limit cycles enclosing an elementary node at the origin when the three-order nilpotent critical point  $O(0,0)$  is a 13-order weak focus.

According to the relations

$$\lambda_1 = \lambda_2 = \lambda_3 = \lambda_4 = \lambda_5 = \lambda_6 = \lambda_7 = \lambda_8 = \lambda_9 = \lambda_{10} = \lambda_{11} = \lambda_{12} = 0, \lambda_{13} \neq 0,$$

we have that

**Theorem 4.1.** *The three-order nilpotent critical point  $O(0,0)$  of system (1.2) is a 13-order weak focus if and only if*

$$\begin{aligned} b_{21} &= -3\mu, \quad a_{12} = -3b_{03}, \quad b_{23} = -a_{32}, \\ a_{14} &= a_{23} - 5b_{05} + 2b_{14}, \quad b_{03} = \mu, \\ a_{32} &= \frac{756 - 142739\mu}{252}, \\ b_{05} &= \frac{189 + 189b_{14} - 35516\mu}{378}, \\ a_{03} &= \frac{945 - 231391\mu}{2646}, \\ b_{14} &= \frac{34020 - 4116a_{23} - 5024095\mu}{1372}, \\ a_{05} &= \frac{-714420 - 594712314\mu + 90016920a_{23}\mu + 102674821303\mu^2}{14002632\mu}, \\ a_{23} &= \frac{359281818 - 36356476968\mu - 2196158788177\mu^2}{42007896}, \\ \mu &\approx \pm 0.00917916. \end{aligned} \tag{4.1}$$

*Proof.* Let

$$\lambda_1 = \lambda_2 = \lambda_3 = \lambda_4 = \lambda_5 = \lambda_6 = \lambda_7 = \lambda_8 = \lambda_9 = \lambda_{10} = \lambda_{11} = 0,$$

we obtain relations of

$$b_{21}, a_{12}, b_{23}, a_{14}, b_{03}, a_{32}, b_{05}, a_{03}, b_{14}, a_{05}, a_{23},$$

Solving the equation  $\lambda_{12} = 0$  about  $\mu$ , we could get four solutions

$$\begin{aligned} A_1 &\approx 0.00358818 - 0.000840459i, \quad A_2 \approx 0.00358818 + 0.000840459i, \\ A_3 &\approx -0.00917916, \quad A_4 \approx 0.00917916. \end{aligned}$$

$\mu \in \mathbb{R}$ , so  $\mu \approx \pm 0.00917916$ , and

$$\begin{aligned} \text{Resultant}[\lambda_{12}, \lambda_{13}, \mu] &= 708862143887297940928622843069693973671321637260998996555 \\ &67040247817324403065696876342634774894135792916822061 \\ &93708806880588535845254301230560000. \end{aligned} \tag{4.2}$$

So  $\lambda_{13} \neq 0$ , the origin of system (1.2) is a 13-order weak focus.



The perturbed system of (1.2) could be written as follows:

$$\begin{aligned} \frac{dx}{dt} &= \delta x + \mu\gamma + \mu x^3 - \mu x^2\gamma + a_{12}x\gamma^2 + a_{03}\gamma^3 + \left(1 - \frac{71275\mu}{378}\right)x^4\gamma + a_{32}x^3\gamma^2 \\ &\quad + a_{23}x^2\gamma^3 + a_{14}x\gamma^4 + a_{05}\gamma^5 - \mu\gamma(x^2 + \gamma^2)^3, \\ \frac{dy}{dt} &= \delta y - 2\mu x^3 + b_{21}x^2\gamma + \mu x\gamma^2 + b_{03}\gamma^3 - 2\left(1 - \frac{71275\mu}{378}\right)x^3\gamma^2 + b_{23}x^2\gamma^3 \\ &\quad + b_{14}x\gamma^4 + b_{05}\gamma^5 + \mu x(x^2 + \gamma^2)^3. \end{aligned} \tag{4.3}$$

When conditions of (4.1) hold, by the relationships  $\lambda_1 = \lambda_2 = \lambda_3 = \lambda_4 = \lambda_5 = \lambda_6 = \lambda_7 = \lambda_8 = \lambda_9 = \lambda_{10} = \lambda_{11} = \lambda_{12} = 0$ , the values of  $b_{21}, a_{12}, b_{23}, a_{14}, b_{03}, a_{32}, b_{05}, a_{03}, b_{14}, a_{05}, a_{23}, \mu$  could be determined. Notice that  $\mu = \pm -0.00917916$  are the simple zeros of  $\lambda_{12} = 0$ . Hence, when conditions in (4.1) hold, we have when  $\mu \approx -0.00917916$

$$\frac{\partial(\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6, \lambda_7, \lambda_8, \lambda_9, \lambda_{10}, \lambda_{11}, \lambda_{12})}{\partial(b_{21}, a_{12}, b_{23}, a_{14}, b_{03}, a_{32}, b_{05}, a_{03}, b_{14}, a_{05}, a_{23}, \mu)} \approx 1.83793 \times 10^{51},$$

when  $\mu \approx 0.00917916$

$$\frac{\partial(\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6, \lambda_7, \lambda_8, \lambda_9, \lambda_{10}, \lambda_{11}, \lambda_{12})}{\partial(b_{21}, a_{12}, b_{23}, a_{14}, b_{03}, a_{32}, b_{05}, a_{03}, b_{14}, a_{05}, a_{23}, \mu)} \approx 4.05781 \times 10^{37}.$$

In fact,

$$\begin{aligned} J &= \frac{\partial(\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5, \lambda_6, \lambda_7, \lambda_8, \lambda_9, \lambda_{10}, \lambda_{11}, \lambda_{12})}{\partial(b_{21}, a_{12}, b_{23}, a_{14}, b_{03}, a_{32}, b_{05}, a_{03}, b_{14}, a_{05}, a_{23})} \\ &= \frac{3125(57367926 - 25174619316\mu + 2196158788177\mu^2)^8}{3553644238836727429387691170820702439358739754227457606557653317975629911417669374\mu^{24} \\ &\quad \times (15338883168 - 6443685685206\mu + 575725672923156\mu^2 + 22094172198662329\mu^3,)} \end{aligned} \tag{4.4}$$

and

$$\text{Resultant}[J, \lambda_{12}, \mu] \neq 0.$$

So when  $\mu \approx \pm 0.00917916, J \neq 0$ .

From the statement mentioned above, Theorem 2.1 follows that

**Theorem 4.2.** *If the three-order nilpotent critical point  $O(0,0)$  of system (1.2) is a 13-order weak focus, for  $0 < \delta \ll 1$ , making a small perturbation to the coefficients of system (1.2), then, for system (4.3), in a small neighborhood of the origin, there exist exactly 13 small amplitude limit cycles enclosing the origin  $O(0,0)$ , which is an elementary node.*

### Appendix A

Detailed recursive MATHEMATICA code to compute the quasi-Lyapunov constants at the origin of system (1.2):

$$c[0, 0] = 0, c[1, 0] = 0, c[0, 1] = 0, c[2, 0] = 0, c[1, 1] = 0, c[0, 2] = 1;$$

when  $k < 0$ , or  $j < 0, c[k, j] = 0$ ;

else

$$\begin{aligned}
 c[k, j] = & \frac{1}{378k(1+s)\mu} (-756\mu c[-8+k, 2+j] - 378j\mu c[-8+k, 2+j] - 756s\mu c[-8+k, 2+j] \\
 & - 378j\mu c[-8+k, 2+j] - 2268\mu c[-6+k, j] - 1134j\mu c[-6+k, j] + 378k\mu c[-6+k, j] \\
 & - 2268s\mu c[-6+k, j] - 1134j\mu c[-6+k, j] + 378ks\mu c[-6+k, j] - 2268\mu c[-4+k, -2+j] \\
 & - 1134j\mu c[-4+k, -2+j] + 1134k\mu c[-4+k, -2+j] - 2268s\mu c[-4+k, -2+j] \\
 & - 1134j\mu c[-4+k, -2+j] + 1134ks\mu c[-4+k, -2+j] + 1512c[-4+k, j] \\
 & - 378kc[-4+k, j] + 1512sc[-4+k, j] + 756j\mu c[-4+k, j] - 378ksc[-4+k, j] \\
 & - 142550j\mu c[-4+k, j] + 71275k\mu c[-4+k, j] - 285100s\mu c[-4+k, j] \\
 & - 142550j\mu c[-4+k, j] + 71275ks\mu c[-4+k, j] + 1512\mu c[-4+k, 2+j] \\
 & + 756j\mu c[-4+k, 2+j] + 1512s\mu c[-4+k, 2+j] + 756j\mu c[-4+k, 2+j] \\
 & + 2268a_{32}c[-3+k, -1+j] + 1512b_{23}c[-3+k, -1+j] - 378b_{23}j c[-3+k, -1+j] \\
 & - 378a_{32}kc[-3+k, -1+j] + 1134a_{32}sc[-3+k, -1+j] + 378b_{23}sc[-3+k, -1+j] \\
 & - 378b_{23}j\mu c[-3+k, -1+j] - 378a_{32}ksc[-3+k, -1+j] - 378b_{21}j c[-3+k, 1+j] \\
 & - 378b_{21}sc[-3+k, 1+j] - 378b_{21}j\mu c[-3+k, 1+j] + 2268\mu c[-3+k, 1+j] \\
 & - 378k\mu c[-3+k, 1+j] + 1134s\mu c[-3+k, 1+j] - 378ks\mu c[-3+k, 1+j] \\
 & - 756\mu c[-2+k, -4+j] - 378j\mu c[-2+k, -4+j] + 1134k\mu c[-2+k, -4+j] \\
 & - 756s\mu c[-2+k, -4+j] - 378j\mu c[-2+k, -4+j] + 1134ks\mu c[-2+k, -4+j] \\
 & + 1512a_{23}c[-2+k, -2+j] + 2268b_{14}c[-2+k, -2+j] - 378b_{14}j c[-2+k, -2+j] \\
 & - 378a_{23}kc[-2+k, -2+j] + 756a_{23}sc[-2+k, -2+j] + 756b_{14}sc[-2+k, -2+j] \\
 & - 378b_{14}j\mu c[-2+k, -2+j] - 378a_{23}ksc[-2+k, -2+j] - 756\mu c[-2+k, j] \\
 & - 378j\mu c[-2+k, j] + 378k\mu c[-2+k, j] - 756s\mu c[-2+k, j] - 378j\mu c[-2+k, j] \\
 & + 378ks\mu c[-2+k, j] + 756a_{14}c[-1+k, -3+j] + 3024b_{05}c[-1+k, -3+j] \\
 & - 378b_{05}j c[-1+k, -3+j] - 378a_{14}kc[-1+k, -3+j] + 378a_{14}sc[-1+k, -3+j] \\
 & + 1134b_{05}sc[-1+k, -3+j] - 378b_{05}j\mu c[-1+k, -3+j] - 378a_{14}ksc[-1+k, -3+j] \\
 & + 756a_{12}c[-1+k, -1+j] + 1512b_{03}c[-1+k, -1+j] - 378b_{03}j c[-1+k, -1+j] \\
 & - 378a_{12}kc[-1+k, -1+j] + 378a_{12}sc[-1+k, -1+j] + 378b_{03}sc[-1+k, -1+j] \\
 & - 378b_{03}j\mu c[-1+k, -1+j] - 378a_{12}ksc[-1+k, -1+j] + 378k\mu c[k, -6+j] \\
 & + 378ks\mu c[k, -6+j] - 378a_{05}kc[k, -4+j] - 378a_{05}ksc[k, -4+j] \\
 & - 378a_{03}kc[k, -2+j] - 378a_{03}ksc[k, -2+j] + 756j\mu c[-4+k, j] - 285100\mu c[-4+k, j]) \\
 \omega_m = & -\frac{1}{378\mu} (378\mu c[-7+m, 1] + 378s\mu c[-7+m, 1] + 756\mu c[-5+m, -1] \\
 & - 378m\mu c[-5+m, -1] + 756s\mu c[-5+m, -1] - 378ms\mu c[-5+m, -1] \\
 & - 1134m\mu c[-3+m, -3] - 1134ms\mu c[-3+m, -3] - 378c[-3+m, -1] \\
 & + 378mc[-3+m, -1] - 71275m\mu c[-3+m, -1] - 756s\mu c[-3+m, 1] \\
 & - 378sc[-3+m, -1] + 378msc[-3+m, -1] + 71275\mu c[-3+m, -1] \\
 & + 71275s\mu c[-3+m, -1] - 71275ms\mu c[-3+m, -1] - 756\mu c[-3+m, 1] \\
 & - 1890a_{32}c[-2+m, -2] - 1890b_{23}c[-2+m, -2] + 378a_{32}mc[-2+m, -2] \\
 & - 756a_{32}sc[-2+m, -2] - 756b_{23}sc[-2+m, -2] + 378a_{32}msc[-2+m, -2] \\
 & - 378b_{21}c[-2+m, 0] - 1890\mu c[-2+m, 0] + 378m\mu c[-2+m, 0] \\
 & + 378ms\mu c[-2+m, 0] - 756\mu c[-1+m, -5] - 1134m\mu c[-1+m, -5] \\
 & - 1134ms\mu c[-1+m, -5] - 1134a_{23}c[-1+m, -3] - 2646b_{14}c[-1+m, -3] \\
 & + 378a_{23}mc[-1+m, -3] - 378a_{23}sc[-1+m, -3] - 1134b_{14}sc[-1+m, -3] \\
 & + 378a_{23}msc[-1+m, -3] - 378m\mu c[-1+m, -1] - 378ms\mu c[-1+m, -1] \\
 & - 3402b_{05}c[m, -4] + 378a_{14}mc[m, -4] - 1512b_{05}sc[m, -4] + 378a_{14}msc[m, -4] \\
 & - 378a_{12}c[m, -2] - 1890b_{03}c[m, -2] + 378a_{12}mc[m, -2] - 756b_{03}sc[m, -2] \\
 & + 378a_{12}msc[m, -2] - 378\mu c[1+m, -7] - 378m\mu c[1+m, -7] - 378s\mu c[1+m, -7] \\
 & - 378ms\mu c[1+m, -7] + 378a_{05}c[1+m, -5] + 378a_{05}mc[1+m, -5] \\
 & + 378a_{05}msc[1+m, -5] + 378a_{03}c[1+m, -3] + 378a_{03}mc[1+m, -3] \\
 & + 378a_{03}msc[1+m, -3] + 378\mu c[1+m, -1] + 378m\mu c[1+m, -1] \\
 & - 756s\mu c[-2+m, 0] - 756s\mu c[-1+m, -5] - 378a_{14}c[m, -4] \\
 & + 378s\mu c[1+m, -1] + 378ms\mu c[1+m, -1] + 378a_{05}sc[1+m, -5] + 378a_{03}sc[1+m, -3]) \\
 \lambda_m = & \frac{\omega_{2m+4}}{2m-4s-1}.
 \end{aligned}$$

#### Acknowledgements

This research was partially supported by the Nature Science Foundation of Shandong Province (Y2007A17, ZR2010AL005) and the National Nature Science Foundation of China (11071222, 11101126).

#### Author details

<sup>1</sup>School of Science, Linyi University, Linyi 276005, Shandong, P.R. China <sup>2</sup>School of Mathematics and Statistics, Henan University of Science and Technology, Luoyang 471003, Henan, P.R. China

#### Authors' contributions

We constructed a concrete system of seventh-degree and investigated the bifurcations of limit cycles from the nilpotent critical point. First of all, the first 13 quasi-Lyapunov constants were deduced. Furthermore, sufficient and necessary conditions in order to have a center were obtained. Moreover, we proved that 13 limit cycles could be bifurcated from its three-order nilpotent critical point. Although this system is very specific, comparing our result with  $N(7) \geq 9$  in [23], we give a lower bound of cyclicity of three-order nilpotent critical point for seventh-degree nilpotent systems. The result of this article is helpful to the 16th Hilbert problem, it enriches the literatures as to the Hilbert's 16th problem. All authors read and approved the manuscript.

#### Competing interests

The authors declare that they have no competing interests.

Received: 21 June 2011 Accepted: 28 February 2012 Published: 28 February 2012

#### References

1. Poincare, H: Memoire sur les courbes definiées par lesequations diffeentielles. *J de Math-ematiques Pures et Appliquees*. **4**, 167–244 (1885)
2. Lyapunov, AM: *Stability of Motion*. Academic Press, New York (1966)
3. Li, J: Hilbert's 16th problem and bifurcation of planar polynomial vector fields. *Int J Bifurcation Chaos*. **13**, 47–106 (2003). doi:10.1142/S0218127403006352
4. Yu, P, Chen, G: Computation of focus values with applications. *Nonlinear Dyn*. **51**, 409–427 (2008). doi:10.1007/s11071-007-9220-7
5. Yu, P: Computation of normal forms via a perturbation technique. *J Sound Vib*. **211**, 19–38 (1998). doi:10.1006/jsvi.1997.1347
6. Kuznetsov, NV, Leonov, GA: Computation of Lyapunov quantities. pp. 1–10. Saint Petersburg, RUSSIA1, (2008) Proceedings of the 6th EUROMECH Nonlinear Dynamics Conference
7. Kuznetsov, NV, Leonov, GA: Lyapunov quantities, limit cycles and strange behavior of trajectories in two-dimensional quadratic systems. *J Vibroeng*. **10**, 460–467 (2008)
8. Moussu, R: Symétrie et forme normale des centres et foyers dégénérés. *Ergodic Theory Dyn Syst*. **2**, 241–251 (1982)
9. Stróżyńska, E, Żołądek, H: The analytic and formal normal form for the nilpotent singularity. *J Diff Equ*. **179**, 479–537 (2002). doi:10.1006/jdeq.2001.4043
10. Takens, F: Singularities of vector fields. *Inst Hautes Études Sci Publ Math*. **43**, 47–100 (1974). doi:10.1007/BF02684366
11. Álvarez, MJ, Gasull, A: Generating limit cycles from a nilpotent critical point via normal forms. *J Math Anal Appl*. **318**, 271–287 (2006). doi:10.1016/j.jmaa.2005.05.064
12. Giacomini, H, Giné, J, Llibre, J: The problem of distinguishing between a center and a focus for nilpotent and degenerate analytic systems. *J Diff Equ*. **227**, 406–426 (2006). doi:10.1016/j.jde.2006.03.012
13. Giacomini, H, Giné, J, Llibre, J: Corrigendum to: The problem of distinguishing between a center and a focus for nilpotent and degenerate analytic systems. *J Diff Equ*. **232**, 702 (2007). doi:10.1016/j.jde.2006.10.004
14. Farr, WW, Li, C, Labouriau, IS, Langford, WF: Degenerate Hopf-bifurcation formulas and Hilbert's 16th problem. *SIAM J Math Anal*. **20**, 13–29 (1989). doi:10.1137/0520002
15. Andronov, AA, Leontovich, EA, Gordon, II, Maier, AG: *Theory of Bifurcations of Dynamical Systems on a Plane*. Wiley, New York (1973)
16. Shi, SL: On the structure of Poincaré-Lyapunov constants for the weak focus of polynomial vector fields. *J Diff Equ*. **52**, 52–57 (1984). doi:10.1016/0022-0396(84)90133-5
17. Álvarez, MJ, Gasull, A: Momodromy and stability for nilpotent critical points, *Internat. J Bifur Chaos*. **15**, 1253–1265 (2005). doi:10.1142/S0218127405012740
18. Chavarriga, J, Giacomini, H, Giné, J, Llibre, J: Local analytic integrability for nilpotent centers. *Ergodic Theory Dyn Syst*. **23**, 417–428 (2003)
19. Gasull, A, Torregrosa, J: A new algorithm for the computation of the Lyapunov constants for some degenerated critical points. *Nonlinear Anal: Proc Illrd World Congress on Nonlinear Anal*. **47**, 4479–4490 (2001)
20. Chavarriga, J, García, I, Giné, J: Integrability of centers perturbed by quasi-homogeneous polynomials. *J Math Anal Appl*. **211**, 268–278 (1997)
21. Liu, Y, Li, J: *Some Classical Problems about Planar Vector Fields* (in chinese). Science press (China), Beijing (2010)
22. Gasull, A, Giné, J: Cyclicity versus Center Problem. *Qual Theory Dyn Syst*. **9**, 101–113 (2010). doi:10.1007/s12346-010-0022-9
23. Andreev, AF, Sadovskii, AP, Tsikalyuk, VA: The center-focus problem for a system with homogeneous nonlinearities in the case of zero eigenvalues of the linear part. *Diff Equ*. **39**, 155–164 (2003). doi:10.1023/A:1025192613518
24. Liu, Y, Li, J: Bifurcation of limit cycles and center problem for a class of cubic nilpotent system. *Int J Bifur Chaos*. **20**, 2579–2584 (2010). doi:10.1142/S0218127410027210

doi:10.1186/1687-1847-2012-20

Cite this article as: Li et al.: Local bifurcation of limit cycles and integrability of a class of nilpotent systems. *Advances in Difference Equations* 2012 **2012**:20.